

Ten-years dataset of poplar inventory in northern Italy

Daniele Rizza¹, Pier Mario Chiarabaglio¹, Domenico Coaloa¹, Alessandro Rocci^{1*}

Received: 15/02/2023 Accepted: 23/05/2023 Published online: 4/10/2023

ABSTRACT The data refer to several poplar plantations located in the plains of northern Italy. The information was collected during the vegetative rest of each year from 1987 to 1996. Dendrometric data were recorded, such as the diameter at breast height, the diameter at five meters height and the planting density, as well as damage caused by biotic and abiotic adversities using a three-level intensity scale. All data is raw, with only total volume and the volume of the first log (up to 5 meters height) calculated using dendrometric equations based on tree diameter and height. The availability of a continuous inventory with annual measurements for 10 years on the same trees in the permanent sample plots has allowed the creation of a particularly important database for the study of growth models and the influence of biotic and abiotic adversities on wood production. This dataset could be used to perform further investigations, such as CO₂ sequestration, to assess the environmental sustainability of the poplar plantations. Furthermore, thanks to this database, it is possible to identify which areas of the northern Italian plains are more suitable for poplar cultivation based on wood biomass production, or to evaluate the impact of pests and diseases with respect to clone and land characteristics.

KEYWORDS: dendrometric data, *Populus* sp., clones, growth modelling, biotic and abiotic adversity data, georeferenced sampling.

Introduction

The growing demand for wood assortments used in timber industry and biomass energy production, combined with efforts to conserve natural forests, has supported a significant expansion of forest plantations. The most commonly used trees belong to fast-growing species such as poplars (Corona et al. 2020).

The genus *Populus* is cultivated to produce sawn timber, veneer, and plywood due to its rapid juvenile growth, high photosynthetic capacity, high growth performance and high wood biomass production (Corona et al. 2020).

The main feature of poplar is its fast-growing nature, which results in short rotations, typically lasting around ten years. This leads to significant inter-annual fluctuations in the geographical distribution of planted areas, requiring frequent assessments that also take into account the growth stage of the trees.

In Italy, poplar cultivation is the most significant source of timber for industrial use, including furniture production, plywood, and other products. Of all the timber processed in Italy, poplar represents half of that which has been grown within the borders of the country.

Poplar cultivation provides several environmental benefits, such as erosion reduction, and is also used for climate change mitigation practices (D'Amico et al. 2021). One of these practices is carbon sequestration, in which trees absorb carbon from the atmosphere and store it in their biomass and the soil, reducing the greenhouse effect and enriching the soil simultaneously.

Additionally, using wood obtained from plantations instead of forests can help reduce pressure on natural fo-

rests and enhance associated environmental benefits.

Forest plantation inventories are valuable data sources for studying the environmental, social, and economic services provided (Nielsen et al. 2014), as they are an important tool for tracking the distribution, characteristics, and time dynamics of plantations (Chiarabaglio et al. 2018).

The aim of this paper is to make available the inventory dataset concerning poplar plantations in northern Italy, so to allow the use in further investigations and analyses.

Methods

Study area


The surveyed land consists of the plains of northern Italy, specifically the Po Valley. Data were collected in the administrative regions of Piemonte, Lombardia, Emilia-Romagna, Veneto and Friuli-Venezia-Giulia. All records are georeferenced with UTM metric coordinates in European Datum 1950, EPSG:4230.

Sampling scheme and data collection

The surveyed land consists of the plains of northern Italy.

2,322 clusters, georeferenced on the three-kilometre UTM grid, are systematically distributed over an area of 4,179,600 hectares. The nine subunits that make up each cluster are in coincidence with the points of a 500 meters square grid (Figure 1). The data refers to surveys during the vegetative rest of each year from 1987 to 1996.

Figure 1 - Clusters and sample units' systematic distribution on the three-kilometer UTM grid 79x82mm (300 x 300 DPI).



The diameter at breast height was measured with a dendrometric caliper, the diameter at 5 m height with a Finnish parabolic caliper. Total height was measured with a Blume Leiss hypsometer. The diameter of the permanent sample plot was measured with a metric tape. The center of permanent sample plot (PSP) was identified on the field

through the interpretation of aerial photos (B/W infrared, scale 1:18,000, period 1978-84) on which their central points have been located using UTM system coordinates ED50 ("European Datum 1950", EPSG:4230).

When the centre of a cluster element falls within an intensive poplar plantation (at least 4 poplar rows covering an area of at least 1,000 m²) a 600 m² circular PSP is defined around such point. Site characteristics are recorded as well as cultivation characteristics, such as poplar clone, spacing, layout and year of planting. All the trees that belong to such sample plot are indelibly marked. Ground surveys begin in spring after budding and when all new plantations have been established. In autumn, at the end of the growing season, a second ground survey is carried out only in six years old and older plantations. Damages caused by biotic and abiotic adversities are surveyed using a scale of three levels of intensity. In the end, the following spring, before a new inventoried cycle takes place, a third ground survey is carried out on six years old and older poplar stands to assess how many, among those existing in the previous autumn, were subsequently felled. Table 1 reports the measures and observations obtained from each survey.

Table 1 - Description of the fields of the dataset

Field	Description	Field type	Unit of measurement
Fuse			
Lat	UTM metric coordinates in the ED50 projection of the center of permanent sample plots (PSP)	Numeric	
Lon			
Cod_ele	The letter refers to the Italian Region code (E= Emilia Romagna, F=Friuli Venezia Giulia, L=Lombardia, P=Piemonte, V=Veneto); the first number specifies the cluster; the second number refers to the element of the cluster	String	
Cod_num	Cod_ele with single tree identification number	String	
h_dendro	Tree height	Numeric	m
h_5m	Tree height at 5 m; less than 5 meters in case of stem breakage or unpruned branches	Numeric	m
Diam	Tree diameter at breast height	Numeric	cm
Diam_5	Tree diameter at 5 meters height	Numeric	cm
Vol (Tree volume)	$Vol = c_1 \cdot D^{c_2} \cdot H^{c_3} \cdot S^{c_4} \cdot a^{c_5}$ c ₁ =0.2535; c ₂ = 2.093, c ₃ = 1.02777; c ₄ =0.0275 c ₅ =0.082; D=Diam; H=h_dendro; S=Planting_density; a=Age	Numeric	m ³
Vol_5 (Tree volume at 5 meters height)	$Vol_5 = \frac{1}{3} \cdot \frac{\pi}{4} \cdot (D^2 + d^2 + D \cdot d) \cdot (h - 1.3) + \frac{\pi}{4} \cdot D^2 \cdot 1.3$ d=Diam_5; h=h_5m	Numeric	m ³
Age	Tree age	Numeric	
Clone	100 P.alba; 200 P.nigra; 300 P.delt. x maxim.; 301 Eridano; 400 P.deltoides; 401 Onda; 402 Lux; 403 Lena; 500 P. x euramericana; 501 I-214; 502 Cima; 503 Pan; 504 Bellini; 505 Cappa Bigliona; 506 Luisa Avanzo; 507 BL Costanzo; 508 Guardi; 510 Type CANADESE; 511 Boccalari; 512 Stella; 513 BRA.NA.GESI; 514 Gattoni; 515 Adige; 520 Type CAROLINIANO; 521 San Martino; 522 Triplo	Numeric	
Growth_stage	1 Tree still in place 2 felled trees (last measurement)	Numeric	

Tree_psp	Number of trees in the permanent sample plot	Numeric	
Planting_density	Plant density	Numeric	m^2
Year_measure		Numeric	
Missing			
Dead		Logic	
Broken			
Replaced			
Planting year		Numeric	
h_1_branch	Height of the first branch	Numeric	m
Mech_damage (mechanical damage)	Damage intensity scale:		
Fire_damage	0 no damage	Numeric	
Hail_damage	1 low		
Frost_damage	2 moderate		
Wind_snow_damage	3 high		
Brown_spot_damage (Marssonina brunnea)			
Virus_damage (Poplar Mosaic Virus)			
Rot_damage (Rosellinia necatrix and others)			
Woolly_aphid (Phloeomyzus passerinii)			
Saperda (Saperda carcharias)			
Rodents			
Necrosis (Discosporium populeum)			

Data access and metadata description

The dataset can be download using the following reference: "Rizza D., Chiarabaglio P.M., Coaloa D., Rocci, A. 2023 - Dataset of ten years of poplar inventory in northern Italy [Data set]." DOI: 10.5281/zenodo.7612569. The file is provided as a 'comma-separated values' file (csv).

Technical validation

The dataset is a collection of dendrometric information, including pest and disease attacks, from intensively cultivated poplar plantations in the plains of northern Italy. It includes data on 33,632 trees measured during a continuous poplar inventory conducted from 1987 to 1996, with annual ground surveys. The data of each tree have been carefully verified before the publication, avoiding misspelling or error among data collected by different operators in different years.

Reuse potential and limits

The ten-year data collection period of this dataset provides a valuable resource for research activities. Sharing this information with the scientific community will allow for the development of growth models, using the georeferenced and temporally placed data to understand the relationships between dendrometric data collected for individual trees and soil and climatic conditions in the sampled areas.

The dataset is particularly useful for modelling the growth of poplar plantations using annual data recorded for each tree, by clone and/or by land characteristics. Additionally, the data could help in assessing the environ-

mental sustainability of the plantations by determining the amount of CO_2 sequestered by the trees.

Acknowledgements

The authors would like to thank the technical staff (Roberto Costanzo, Michele Sciré and Tiziano Collot) who made it possible to collect the data and the colleague Gianfranco Lapietra who set up and started the inventory.

References

Chiarabaglio P.M., Bergante S., Scirè M., Coaloa D. 2018 - *Low cost poplar inventory in the plain of Piemonte (Italy)*. Annals of Silvicultural Research 42 (1): 39-42.

Corona P., Chianucci F., Marcelli A., Gianelle D., Fattorini L., Grotti M., Puletti N., Mattioli W. 2020 - *Probabilistic sampling and estimation for large-scale assessment of poplar plantations in northern Italy*. European Journal of Forest Research 139 (6): 981-988.

D'Amico G., Francini S., Giannetti F., Vangi E., Travagliani D., Chianucci F., Chirici G. 2021 - *A deep learning approach for automatic mapping of poplar plantations using sentinel-2 imagery*. GIScience and Remote Sensing 58 (8): 1352-1368.

Nielsen A.B., Östberg J., Delshammar T. 2014 - *Review of urban tree inventory methods used to collect data at single-tree level*. Arboriculture & Urban Forestry 40 (2): 96-111.