

Salicylic acid alleviates methyl viologen induced oxidative stress through transcriptional modulation of antioxidant genes in *Zea mays L*

Pranjali Yadava^{1*}, Nepolean Thirunavukkarasu², Palminder Kaur¹, Kaliyugam Shiriga², Ishwar Singh^{1*}

¹ICAR - Indian Institute of Maize Research, Pusa Campus, New Delhi 110 012, India

²Division of Genetics, ICAR- Indian Agricultural Research Institute, Pusa Campus, New Delhi 110 012, India

*Corresponding author: E-mail: isingh.dmr@gmail.com; pranjali.yadava@icar.gov.in

Abstract

Salicylic acid (SA) is an important growth regulator that participates in both biotic as well as abiotic stress response in plants. Both-biotic and abiotic stresses are characterized by elevated cellular reactive oxygen species (ROS). We examined the biochemical and molecular effects of SA on the ROS scavenging system in maize. Pre-treatment of maize plants with SA prior to stress, reduced electrolyte leakage, enhanced protein accumulation, improved root biomass and grain yield. SA application also led to a general enhancement in the biochemical activities of three major ROS scavenging antioxidant enzymes, viz. superoxide dismutase (SOD), catalase (CAT) and peroxidases (POX). However, corresponding transcripts of various antioxidant isozymes depicted a dynamic response for different SA doses. Under stress, pre-treatment with 0.5 mM SA led to enhanced expression of all the Sod genes analyzed while higher SA concentration repressed many Sod genes. *Cat 1* showed clear dose-dependent repression in response to graded concentrations of SA, while *Cat 3* showed inverse dose-dependent activation in response to graded concentrations of SA. *Apx 1* was found to be up-regulated at 0.5 mM SA, while higher doses of SA, i.e. 1.0 and 1.5 mM led to its repression. All the three concentrations of SA repressed *Apx 2* gene. Taken together, our results indicate that SA has the potential to alleviate oxidative stress in maize through biochemical and transcriptional modulation of ROS scavenging pathway. Also, the concentration of SA has a great bearing in its alleviating role, with lower concentrations (viz 0.5 mM) being more desirable.

Keywords: salicylic acid, reactive oxygen species, antioxidant genes, oxidative stress, maize

Introduction

Salicylic acid (SA) has long been known for its role in the responses to biotic stresses in plants, especially, its ability to induce systemic acquired resistance against viral diseases (Gaffney et al, 1993). It participates in diverse signalling pathways and has cross-talk with various plant growth regulators and hormones. Recent studies have suggested that it can also modulate pathways leading to adaptation to abiotic stresses in plants (Kang et al, 2014). One of the common consequence of pathogen attack as well as that of incidence of abiotic stress is the rapid burst of reactive oxygen species (ROS). Persistent accumulation of high levels of ROS is detrimental for plant survival. Therefore, plants have evolved mechanisms to scavenge harmful ROS, in which, antioxidant enzymes like superoxide dismutase (SOD), catalase (CAT) and peroxidases (POX) play a major role. There is an increasing interest in the interactive role between SA and ROS (Dat et al, 2007). SA occurs naturally in plants in very low amounts and it is known to be involved in transpiration, stomatal closure, nutrient uptake, protein synthesis, inhibition of ethylene biosynthesis, plant growth, thermogenesis, flower induction, photosynthesis, enzyme activities and other

important regulatory physiological processes (Yusuf et al, 2013). There have been conflicting views on the effect of SA on ROS-mediated oxidative damage. Initial evidence indicated that SA was a potent inhibitor of heme-containing enzymes, viz. CAT and POX, and an inducer of hydrogen peroxide (H_2O_2) (Harfouche et al, 2008), thus further exacerbating accumulation of ROS under stress conditions. SA was also proposed as both a potent inducer of the NADPH-oxidase and an inhibitor of alternative oxidase, thus capable of indirect regulation of the redox status of the plant. To reconcile a considerable body of contradictory views on the question, whether SA is upstream of ROS or vice versa, several researchers proposed that SA and H_2O_2 form a self-amplifying feedback loop (Vlot et al, 2009).

There have been several studies, where pre-treatment of plants with SA has been shown to confer abiotic stress tolerance of various degrees. Cucumber and tobacco plants pre-treated with SA exhibited enhanced protection to oxidative stress induced by methyl viologen (MV) application at 3 days after the SA pre-treatment (Strobel and Kuc, 1995). Similarly, pre-treatment of barley seedlings with SA in the dark via the transpiration stream fully blocked the

subsequent light-induced inhibitory effect of MV on photosynthesis and also decreased the MV-induced production of H_2O_2 , lipid peroxidation, and electrolyte leakage (Ananieva et al, 2002), while inducing the activities of certain antioxidant enzymes (Ananieva et al, 2004). These results were also confirmed using SA-deficient NahG transgenic rice plants, which showed great sensitivity to MV. These plants also had lower glutathione content, which may explain the greater sensitivity of these plants to oxidative damage and also suggests that SA may regulate glutathione synthesis in rice (Kusumi et al, 2006).

Although exogenous SA mediated perturbation in cellular redox state under stress has been studied in many plants, its mechanism of action, especially its effect on transcriptional dynamics is unexplored. This study was undertaken to determine the physiological and biochemical changes in maize plants treated with SA during MV-induced oxidative stress, and to investigate the role of this plant growth regulator in altering the transcriptional dynamics of the antioxidant genes. Elucidating the mechanisms of plant growth regulator mediated stress adaptation in plants would pave way for developing appropriate spray based technologies or transgenic crops that are resilient to environmental stresses.

Materials and Methods

Plant growth and treatments

This experiment was conducted at the Maize Physiology Field Research Facility, ICAR- Indian Agricultural Research Institute, New Delhi, India (Latitude: 28°03'23"N, Longitude: 77°09'27"E, Altitude: 228.61 masl) during July-October 2012. The mean maximum and minimum temperatures during the period of study were 39.1 and 13.9°C, respectively, while total rainfall was 481.8 mm. HQPM 1, a single cross (HKI 193-1 x HKI 163) quality protein maize hybrid was grown in micro-plots of 3 rows with 2 m row length with 10 plants in each row. The crop was irrigated only once just after sowing and remained completely rainfed thereafter. Recommended fertilization and management practices were followed. Plants were sprayed with different doses of SA @ 0.5, 1.0 and 1.5 mM at the flowering stage. All the spray formulations contained Tween 20 (0.1%) as surfactant. The control plants were mock sprayed with only Tween 20 dissolved in distilled water. Artificial oxidative stress was induced in the plants by spraying 200 μ M MV. Leaves were harvested from five different plants per treatment at two stages i.e. after 48 h from application of SA (post treatment) and after 48 h from MV application (post stress); snap freezed and stored at -70°C.

Root growth and yield

At harvest, the data on total root biomass and root length were recorded and expressed as g per plant and cm per plant, respectively. The cobs were harvested from the SA treated and untreated plants

at physiological maturity (30 % grain moisture). The final grain yield at 14% grain moisture was calculated as g per plant.

Membrane stability

Five leaf discs weighing 100 mg each from five different plants per treatment were washed, dried and placed in a glass test tube containing 20 ml deionized water in replicates of three for each treatment. The samples were kept for 24 hrs at room temperature and initial conductance values (E1) were recorded with conductivity meter. The content was then boiled for 30 min in a boiling water bath (100°C), cooled to room temperature and final conductivity values (E2) were determined. Cell membrane injury was represented in terms of total electrolyte leakage (%) and calculated by following formula:

$$\text{Electrolyte Leakage (\%)} = \frac{\text{Conductivity before boiling (E1)}}{\text{Conductivity after boiling (E2)}} \times 100$$

Total soluble proteins and anti-oxidant enzyme assays

Leaf samples (1 g) were homogenized in 10 ml ice-cold 50 mM potassium phosphate buffer (pH 7) containing 100 mg g⁻¹ of polyvinyl pyrrolidone (PVP) in mortar and pestle and centrifuged at 10,000 rpm for 30 min at 4°C. The supernatant was then collected and stored at -20°C. The supernatant collected was used for different antioxidant enzyme assays after estimating total protein content (Bradford, 1976) using bovine serum albumin as standard.

Superoxide dismutase (SOD) was assayed according to the method of Kakkar et al (1984). The assay mixture contained 1.2 ml of sodium pyrophosphate buffer (0.025 M, pH 8.3), 0.1 ml of phenazine methosulphate (PMS) (186 μ M), 0.3 ml of nitro blue tetrazolium (NBT) (300 μ M), 0.2 ml of the enzyme extract and water in a total volume of 2.8 ml. The reaction was initiated by the addition of 0.2 ml of NADH. The mixture was incubated at 30°C for 90 s and arrested by addition of 1.0 ml of glacial acetic acid. The reaction mixture was then shaken with 4.0 ml of n-butanol, allowed to stand for 10 min and centrifuged. The intensity of the chromogen in the butanol layer was measured at 560 nm in a spectrophotometer. One unit of enzyme activity is defined as the amount of enzyme that gave 50% inhibition of NBT reduction in one minute. Catalase (CAT) activity was assayed using the method of Aebi (1983). The assay solution contained 50 mM potassium phosphate buffer (pH 7) and 30 mM H_2O_2 and 50 μ l enzyme extract. The activity was estimated by monitoring the decrease in absorbance of H_2O_2 within 30 s at 240 nm. Unit of enzyme activity was taken as the change in absorbance per minute at 240 nm. The non-specific peroxidases (POX) activity was assayed using the method of Reddy et al (1995). To 3.0 ml of pyrogallol solution (0.05 M), 0.1 ml of the enzyme extract was added and the spectrophotometer was adjusted to read zero at

Table 1 - List of primer pairs used for the qRT-PCR amplification of 11 antioxidant genes in maize.

S No.	Gene name	Primer Sequence
1	Zm Sod2	F-GATCATTGGCCGAGCTGTTGTTG R-TCTATGGTGTGCTGGAGGTT
2	Zm Sod4	F-AAACAAATGGGTCGCTGGACTCG R-ACAGTTCAGCTCATGGGTGCTTA
3	Zm Sod9(4A)	F-CGTGTTGCTGTGGATCATTGA R-TGGTGGCTACAGGTGCATAATGA
4	Zm Cu-Zn Sod (chl)	F-TGTTGCAAATGCTGAGGGCATAGC R-CCAACAACACACATGCCAGTCTT
5	Zm Fe Sod	F-AGCACAGGTCTGGAAACCATCACTT R-ACAGCTGAAGGCTGAGCGGATAA
6	Zm Mn Sod	F-TGTTGACCTGCTGGACCAAGTGT R-ACTACGAGCAGCAGAAAGTGGAGT
7	Zm Apx1	F-TGCTGCCTGTATGGAATCATGTGG R-TACCAACAAACAGCTTAAGGGCTCA
8	Zm Apx2	F-TGTTGACAATAACGCAGCGGACG R-ACACTCTCTGGTTGTTACGCCT
9	Zm Apx 8	F-AGATGCAGATGACCAGGAAGCGT R-TCGGAAGGTGAAGGAGCTTGTCA
10	Zm Cat1	F-ACCTGAAGCCGAGCATGTAAGGAT R-ATAATCGACCCACCGACCACATCAGCA
11	Zm Cat3	F-CCGGCTCAACATGAAGGAAACAT R-TTCTCTTGTCTGGCGACGACAT

430 nm. To the test cuvette, 0.5 ml of H_2O_2 (1%) was added and mixed. The change in absorbance was recorded every 30 s up to 3 min in a spectrophotometer. One unit of peroxidase was defined as the change in absorbance per min at 430 nm.

Total RNA isolation and cDNA preparation

Total RNA was extracted from pools of three plants for each experimental sample using the RNeasy mini kit (Qiagen, Hilden, North Rhine- Westphalia, Germany) followed by treatment with DNase I to remove any genomic DNA contamination. Concentrations were determined with a NanoDrop spectrophotometer (Thermo scientific NanoDrop 1000). The cDNA was synthesized using Superscript III Kit (Invitrogen) from 1 μ g of total RNA according to the manufacturer's protocol. The reverse transcription reaction was carried out at 44°C for 60 min followed by 92°C for 10 min. The reaction without template was included as control.

Quantitative Real Time -PCR Analysis

The sequences of the corresponding maize genes of the ROS scavenging system were obtained from the GenBank. Using these sequences, 11 sets of primers were designed using IDT PrimerQuest for further gene expression analysis (Table 1). For the quantitative real-time PCR (qRT-PCR), 1 μ L cDNA (10 ng) was used in 25 μ L reactions using Hot start-IT SYBR green qPCR master mix (Affymetrix) according to the manufacturer's instructions. Amplification was performed on the Mx3005P multiplex quantitative PCR system (Stratagene). For each sample, at least three replications were performed in one experiment. The threshold cycles (CT) of each test target were averaged for triplicate reactions. 18S rRNA was used as internal control (Schmittgen and Livak, 2008). The gene expression values were normalized

by subtracting the mean reference gene (18S rRNA) CT value from individual CT values of corresponding target genes (Δ CT). The fold change value was calculated using the expression, where $\Delta\Delta$ CT represents difference between the Δ CT condition of interest (i.e. gene expression in SA treated plants after MV stress) and Δ CT control (i.e. gene expression in mock treated plants after MV stress).

Results and Discussion

Root growth and yield

The SA treated plants were monitored till the physiological maturity stage. Significantly higher root biomass and grain yield were observed in SA treated plants over the control. Although marginal increase in root length was also observed in SA treated plants over control but this increase was non-significant. The most effective concentration of SA was 1 mM, which improved root biomass and grain yield by 45.7 and 32.9% over untreated plants (Table 2), indicating that SA is capable of alleviating harmful effects of oxidative stress in maize.

Previous studies corroborate that lower concentrations of SA (viz. 0.1 mM to 0.5 mM) are most optimal in imparting stress tolerance in plants with low levels of resident SA, maize being one of them (Janda et al, 1999; Ding et al, 2002; Kang and Saltveit, 2002; Kang et al, 2003; Nemeth et al, 2002; Tasgin et al, 2003; He et al, 2005; Shi et al, 2006). Many previous studies demonstrate that SA could be a potential compound for the reduction of abiotic stress sensitivity of the crops against a range of abiotic stresses, like ozone (Rao and Davis 1999; Koch et al, 2000), salt and osmotic (Borsani et al, 2001; Molina et al, 2002), UV-B (Surplus et al, 1998; Nawrath et al, 2002), drought (Nemeth et al, 2002; Munne-Bosch and Penuelas, 2003), paraquat (Yang et al, 2004), heat (Clarke et al, 2004; He et al, 2005; Shi et al, 2006), cold (Janda et al, 1999; Kang and Saltveit, 2002; Kang et al, 2003; Tasgin et al, 2003; Scott et al, 2004) and metal stress (Metwally et al, 2003; Yang et al, 2003; Pal et al, 2005). Application of ascorbic acid has also been reported to have similar effects on root biomass and grain yield in maize under MV induced oxidative stress (Yadava et al, 2013).

Membrane stability

In order to understand the effect of SA in imparting stability to the cell membrane, we compared total electrolyte leakage in SA treated plants under stress with that of untreated plants under stress. SA treatment led to significant reduction in electrolyte leakage under stress. Under stressed condition, each of the SA treatment, i.e. 0.5, 1.0, and 1.5 mM reduced the electrolyte leakage by 2.7, 4.4 and 3.3%, respectively over the mock treated control. Interestingly, SA was capable of reducing ion leakage under unstressed conditions also, indicating its general role in imparting stability to the membrane. A reduction of 10.4, 15.4, and 11.77% in electrolyte leakage was

Table 2 - Root biomass, root length and grain yield per plant at maturity for different treatments of SA under stressed conditions. Each value represents mean of three independent replications \pm SE.

SA Conc. (mM)	Root biomass (g plant ⁻¹)	Root length (cm plant ⁻¹)	Grain weight (g plant ⁻¹)
0.0	12.39 \pm 1.45	27.36 \pm 1.21	38.42 \pm 2.87
0.5	15.32 \pm 2.31	28.12 \pm 2.07	43.42 \pm 1.94
1.0	18.05 \pm 1.79	29.37 \pm 1.94	51.07 \pm 3.17
1.5	16.21 \pm 1.03	28.61 \pm 1.78	46.37 \pm 2.86

observed in response to 0.5, 1.0, and 1.5 mM SA under unstressed conditions (Figure 1A).

In another study also, SA application was found to reduce electrolyte leakage under different abiotic stresses. In tomato plants, application of 0.1 mM SA led to 44% and 32% reduction in electrolyte leakage in plants subjected to 150 mM NaCl and 200 mM NaCl, respectively, compared to the untreated plants (Stevens et al, 2006). Supporting evidences for SA mediated reduction in electrolyte leakage have also been reported in maize, rice and cucumber hypocotyls under chilling stress (Kang and Saltviel, 2001). Maintaining integrity of cellular membranes under stress conditions is considered an integral part of the tolerance mechanism.

Total soluble proteins

In plants, oxidative stress generally leads to enhanced synthesis and accumulation of certain proteins, especially those related to ROS scavenging and chaperone functions (Horvath et al, 2007). We also found enhanced total soluble proteins in all the plants sprayed with MV in comparison to unstressed plants. Next, we were interested to know whether application of SA had any effect on accumulation of total soluble proteins. In unstressed conditions, 48 h after SA treatment, total soluble protein content of the samples showed an inverse dose dependent response. The protein content increased by 17.8, 14.4, and 12.5% in response to 0.5, 1.0, and 1.5 mM SA, with 0.5 mM concentration being most effective in increasing the protein content. Further, we studied the effect of SA treatment on protein content under stress conditions. Under stress, different doses of SA were found to be capable of enhancing the protein content by 8.3, 3.6, and 1.2% in comparison to the mock control. Here also, inverse dose dependence as seen for unstressed conditions was observed, with 0.5 mM concentration being statistically most significant (Figure 1B).

Since, SA acts as a signalling molecule; it has potential to up-regulate several genes and their products. We also observed enhanced accumulation of total soluble proteins in response to SA application, especially so with lower concentration of 0.5 mM. Evidence for SA mediated accumulation of several classes of proteins, viz. topoisomerases, elongation factors, transcriptionally active long terminal repeat retrotransposon families, heat-shock proteins (Hsp70, Hsp17.6, Hsc70), cyclophilins, dehydrins,

pathogenesis related proteins, glycine-betaine etc. have been provided in various studies (Horvath et al, 2007). Also, physiologically lower concentrations, viz 0.1 to 0.5 mM have been reported to be most effective in various studies.

Antioxidant enzyme activities

Plants exhibit a fine homeostasis between production and scavenging of ROS. The ability of the plant to rapidly quench extra ROS production during stress determines its tolerance capacity (Gill and Tuteja, 2010). SOD, CAT and POX are the important ROS scavenging enzymes in plants. Therefore, we assayed the biochemical activities of these enzymes in response to different doses of SA before and after application of oxidative stress (Figure 2). We observed that MV can drastically reduce the specific activity of SOD. There was 54.4% reduction in SOD activity in mock sprayed plants after application of MV. During unstressed conditions, lower doses of SA application




Figure 1 - Total electrolyte leakage (a) and total soluble protein (b) in response to different concentrations of SA under control and stressed conditions in maize. Each value represents mean of three independent replications \pm S.E, asterisk indicates significant difference compared to the control post stress at $P < 0.05$.




Figure 2 - Specific activities of superoxide dismutase (a), catalase (b) and peroxidases (c) in response to different concentrations of SA under control and stressed conditions in maize. Each value represents mean of three independent replications \pm S.E, asterisk indicates significant difference compared to the control post stress at $P < 0.05$.

(0.5 mM and 1.0 mM) led to reduced SOD activities while the higher dose of SA (1.5 mM) increased SOD activity marginally by 9.0%. However, under stressed conditions, SA was found to have considerable effect in enhancing SOD activity. An increase of 62.6, 104.8, and 144.8% in SOD activity was observed in stressed plants pre-treated with 0.5, 1.0, and 1.5 mM SA respectively, as compared to the mock treated plants. In contrast to SOD, a general enhancement in CAT activity was observed upon induction of oxidative stress. SA was found to further increase CAT activity under stress. Under stress, CAT activity increased by 7.3, 49.3, and 41.3% in response to 0.5, 1.0, and 1.5 mM SA, with later two concentrations having most significant effect. Like SOD, non-specific POX activity also showed a general reduction under MV induced oxidative stress. Once again, confirming its role in alleviating oxidative stress, SA was found to enhance POX activity. Application of all the concentrations of SA increased POX activity in both stressed as well as unstressed conditions. Under stressed conditions, pre-treatment of 0.5, 1.0, and 1.5 mM SA increased

POX activity by 5.3, 8.9, and 7.9% respectively as compared to mock sprayed plants. Thus, SA application led to enhanced activities of all the three ROS scavenging enzymes (SOD, CAT, and POX) under stress, highlighting its positive role in alleviating oxidative stress.

In our experiment, SA application led to enhanced activities of all the three scavengers, i.e. SOD, CAT, and POX. In many previous studies, SA mediated alteration in various antioxidant enzyme activities under stress has been reported. However, there exist considerable inconsistencies and variations in changes in enzyme activities. The response has been largely dependent on genotypes in question, the dose of SA, time of application and nature of abiotic stress. For example in a recent study in wheat, treatment with SA alone increased all the antioxidant activities in Gerek-79 genotype (Multu and Atici, 2013). However, the same treatments with SA decreased the activities of CAT and SOD in Bezostaya genotype, while increased that of POX. On the other hand, salt applications alone increased the activities of POX and SOD, while decreased that of CAT in Bezostaya genotype. The same salt applications increased the POX activity, while decreased the activities of CAT and SOD in Gerek-79 genotype. In plants under saline conditions, except 0.75 M NaCl, treatments with SA increased the activities of CAT, POX, and SOD in both the varieties compared with plants applied with salt alone. In plants applied with 0.75 M NaCl, however, treatments with SA decreased CAT activity. In another study, incubation of *Arabidopsis* leaves with SA enhanced SOD activity in a dose-dependent manner, while treatment of leaves with 5 mM SA decreased CAT by 33% compared with control leaves. In the same study, concentrations of 0 to 5 mM SA had no major effect on POX activity (Rao et al, 1997). We observed that MV application can drastically reduce SOD activity. Loss of ROS scavenging capacity due to reduction in SOD activity might be a major factor contributing to lethal effects of MV induced oxidative burst. In contrast to SOD, a general enhancement in CAT activity was observed upon induction of oxidative stress. This may be due to the crucial need of the plant to expeditiously remove harmful H₂O₂ produced due to MV induced severe oxidative stress.

Antioxidant gene expression analysis

In order to understand more about the molecular mechanisms that may contribute to the variation of the ROS scavenging enzymes, we studied the transcriptional dynamics of the corresponding genes. Though biochemically indistinguishable, some of these scavenging enzymes have a number of isozymes. For example, SODs are classified into three types: Mn-, Fe-, and Cu/Zn-SOD, depending on the metal found in the active site. In the higher plants, SODs are found in the cytosol, plastids, and mitochondria. In maize, different SOD isozymes are reported: four Cu/Zn cytosolic isozymes (SOD-2, SOD-4, SOD-4A,




Figure 3 - Fold change in transcript abundance of various *Sod* (a), *Cat* (b), and *Apx* (c) genes in SA pre-treated plants under stress as compared to control plants under stress. Each value represents mean of three independent replications \pm S.D. Negative values show gene repression.

and SOD-5), four mitochondrial-associated Mn SODs (SOD-3.1, SOD-3.2, SOD-3.3, and SOD-3.4), and one Cu/Zn chloroplast-associated isozyme (Baum and Scandalios, 1979; Scandalios, 1997; Guan and Scandalios, 1998). Also, in maize genome, there exist multiple gene homologs coding for one group of scavenging enzymes. We designed DNA primers that were capable of distinguishing various homologs of these genes (Table 1).

Figure 3 depicts fold change of transcript accumulation in SA pre-treated plants after 48 hrs of MV treatment as compared to mock treated controls. Under stress, pre-treatment with 0.5 mM SA led to enhanced accumulation of transcripts of all the *Sod* genes analyzed. The fold increase ranged from 1.4 times for *Sod* 4 to 3.2 times for *Fe Sod* transcript. The positive fold change for other *Sod* transcripts was 2.3, 2.6, 1.5, and 1.8 times for *Cu/Zn Sod*, *Mn Sod*, *Sod* 2 and *Sod* 9 respectively. When the concentration of SA was doubled to 1.0 mM, fold increase in transcript accumulation for various *Sod* genes was decreased as compared to that of 0.5 mM dose, with *Sod* 2 showing repression. At 1.0 mM SA concentration *Cu/Zn Sod*, *Fe Sod*, *Mn Sod*, *Sod* 4 and *Sod* 9 genes were up-regulated by 1.3, 1.7, 1.4, and 1.1 times respectively as compared to the mock control. On the other hand, *Sod* 2 was repressed by 1.0 fold. When, the concentration of SA spray was further enhanced to 1.5 mM, even more number of *Sod* genes showed repression. In this condition, *Mn Sod* exhibited a repression of 4.9 times, while *Sod* 4 and *Sod* 9 showed 2.5 and 1.6 fold repression respectively. Thus, a dynamic response of transcript accumulation

of various *Sod* genes was observed, which was also affected by the concentration of SA, with 0.5 mM SA being most effective in uniformly up-regulating various isozymes of *Sod*. At the protein level also, SA mediated enhancement of SOD enzyme activity under stress was most significant for 0.5 mM concentration. This also correlates with significantly higher total protein accumulation observed in samples pre-treated with 0.5 mM SA concentration.

We analyzed two *Cat* genes, *Cat 1* and *Cat 3* and both these genes showed contrasting transcriptional profile. *Cat 1* showed clear dose dependent repression in response to graded concentrations of SA, while *Cat 3* showed inverse dose dependent activation in response to graded concentrations of SA. Also, sensitivity of *Cat* genes to SA was seen to be very high for both activation and repression. *Cat 1* expression reduced by 5.0, 15.7, and 34.1 fold respectively in response to 0.5, 1.0 and 1.5 mM SA. On the other hand, *Cat 3* expression increased by 22.3, 4.6, and 2.6 folds respectively in response to 0.5, 1.0, and 1.5 mM SA, with 0.5 mM SA once again being most effective. Transcriptional profile of the three *Apx* genes analyzed exhibited dynamic response. *Apx 1* was found to be up-regulated by 1.6 fold at 0.5 mM SA, while higher doses of SA, i.e. 1.0 and 1.5 mM led to repression of *Apx 1* by 2.5 and 1.4 times respectively. All the three concentrations of SA repressed *Apx 2* gene. *Apx 2* transcript accumulation was reduced by 1.2, 4.6, and 2.2 times respectively in stressed plants pre-treated with 0.5, 1.0, and 1.5 mM SA. On the other hand, *Apx 8* transcript abundance increased 2.5 and 1.2 folds in response to 0.5 mM and 1.5 mM SA, while it decreased 1.4 fold in response to 1.0 mM SA.

The dynamic transcriptional profile of various genes of the ROS scavenging pathway in response to various doses of SA suggest that different ROS genes are differentially regulated by SA and may also likely to form a feedback system. For example, originally, H_2O_2 was proposed to function downstream of SA on the basis of evidence that high levels of SA can bind and inhibit H_2O_2 -removing enzymes, such as CAT and APX (Durner and Klessig, 1996; Channongpol et al, 1998). Furthermore, APX is post-transcriptionally suppressed by SA (Mittler et al, 1998; Yuan and Lin, 2004), and CAT is downregulated at the level of steady-state mRNA (Dorey et al, 1998). Later, it was found that increased H_2O_2 levels also act upstream of SA to induce endogenous SA (Leon et al, 1995). Thus, SA and H_2O_2 compose a positive feedback loop. Moreover, endogenous SA concentration of the plant can directly affect accumulation of ROS. This has been shown in SA deficient transgenics and mutants in Arabidopsis. Transgenic Arabidopsis expressing a bacterial salicylate hydroxylase NahG, and Arabidopsis mutant sid2 (SA induction deficient) and eds5 (enhances disease susceptibility), which have a very low level of SA (below 1 μ g g⁻¹ fresh weight), accumulate a high level ROS and therefore prone to

more severe damages from abiotic stress (Lederer and Böger, 2003; Nawrath and Metraux, 1999; Rao and Davis, 1999; Nawrath et al, 2002; Clarke et al, 2004). Recent studies have also shown that SA is capable of inducing promoters and transcription factors (Van der Does et al, 2013).

In conclusion, our results suggest that SA have potential to alleviate oxidative stress and biochemical and transcriptional modulation of ROS scavenging pathway might be a key mechanism behind its protective role. Also, the concentration of SA is important in its alleviating role, with lower concentrations (viz. 0.5 - 1.00 mM) being more favourable. The increase in SOD, CAT, and POX activities and associated transcriptional modulation with the application of SA, strengthen the conjecture that SA might have neutralized the harmful ROS by not only acting as antioxidant molecule itself but also as a signalling molecule causing up-regulation of expression of genes and specific factors related to synthesis of antioxidant enzymes. The results of the investigation confirm the importance of exogenously applied SA in conferring protection against oxidative stress in maize through modulation of antioxidant pathway.

Acknowledgements

Financial support from Indian Council of Agricultural Research (ICAR) funded Network Project on Transgenics in Crops- Functional genomics for drought tolerance in maize (NPTC 3015) is thankfully acknowledged.

References

Aebi H, 1983. Catalase. In: Methods of enzymatic analysis. Bergmeyer H ed. Verlag Chemical, Weinheim

Ananieva EA, Alexieva VS, Popova LP, 2002. Treatment with salicylic acid decreases the effects of paraquat on photosynthesis. *J Plant Physiol* 159: 685-693

Ananieva EA, Christov KN, Popova LP, 2004. Exogenous treatment with salicylic acid leads to increased antioxidant capacity in leaves of barley plants exposed to paraquat. *J Plant Physiol* 161: 319-328

Baum JA, Scandalios JG, 1979. Developmental expression and intracellular localization of superoxide dismutases in maize. *Differentiation* 13: 133-140

Borsani O, Valpuesta V, Botella MA, 2001. Evidence for a role of salicylic acid in the oxidative damage generated by NaCl and osmotic stress in *Arabidopsis* seedlings. *Plant Physiol* 126: 1024-1030

Bradford MM, 1976. Rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72, 248-254

Chamnongpol S, Willekens H, Moeder W, Langebar-

tels C, Sandermann H, Van Montagu A, Inze D, Van Camp W, 1998. Defense activation and enhanced pathogen tolerance induced by H₂O₂ in transgenic tobacco. *Proc Natl Acad Sci USA* 95: 5818-582

Clarke SM, Mur LAJ, Wood JE, Scott IM, 2004. Salicylic acid dependent signaling promotes basal thermotolerance but is not essential for acquired thermotolerance in *Arabidopsis thaliana*. *Plant J* 38: 432-447

Dat JF, Capelli N, Breusegem FV, 2007. Salicylic acid: a plant hormone, pp. 247-276. In: The interplay between salicylic acid and reactive oxygen species during cell death in plants. Hayat and Ahmad eds. Springer Netherlands

Ding CK, Wang CY, Gross KC, Smith DL, 2002. Jasmonate and salicylate induce the expression of pathogenesis-related-protein genes and increase resistance to chilling injury in tomato fruit. *Planta* 214: 895-901

Dorey S, Baillieul F, Saindrenan P, Fritig B, Kauffmann S, 1998. Tobacco class I and class II catalases are differentially expressed during elicitor-induced hypersensitive cell death and localized acquired resistance. *Mol Plant-Microbe Interact* 11: 1102-1109

Durner J, Klessig DF, 1996. Salicylic acid is a modulator of tobacco and mammalian catalases. *J Biol Chem* 271: 28492-28501

Gaffney T, Friedrich L, Vernooy B, Negrotto D, Nye G, Uknnes S, Ward E, Kessmann H, Ryals J, 1993. Requirement of salicylic acid for the induction of systemic acquired resistance. *Science* 261: 754-756

Gill SS, Tuteja, N, 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol Biochem* 48(12): 909-930

Guan L, Scandalios JG, 1998. Two structurally similar maize cytosolic superoxide dismutase genes, *sod4* and *sod4a*, respond differentially to abscisic acid and high osmoticum. *Plant Physiol* 117(1): 217-224

Harfouche AL, Rugini E, Mencarelli F, Botondi R, Muleo R, 2008. Salicylic acid induces H₂O₂ production and endochitinase gene expression but not ethylene biosynthesis in *Castanea sativa* in vitro model system. *J Plant Physiol* 165(7): 734-44

He YL, Liu YL, Cao WX, Huai MF, Xu BG, Huang BR, 2005. Effects of salicylic acid on heat tolerance associated with antioxidant metabolism in Kentucky bluegrass. *Crop Sci* 45: 988-995

Horváth E, Szalai G, Janda T, 2007. Induction of abiotic stress tolerance by salicylic acid signalling. *J Plant Growth Regul* 26: 290-300

Janda T, Szalai G, Tari I, Paldi E, 1999. Hydroponic treatment with salicylic acid decreases the effects of chilling injury in maize (*Zea mays* L) plants. *Planta* 208: 175-180

Kakkar P, Dos B, Viswnathan PN, 1984. A modified spectrophotometric assay of superoxide dismutase. *Indian J Biochem* 21: 130-132

Kang G, Li G, Guo T, 2014. Molecular mechanism of salicylic acid-induced abiotic stress tolerance in higher plants. *Acta Physiol Plant* 36: 931-943

Kang GZ, Wang ZX, Sun GC, 2003. Participation of H₂O₂ in enhancement of cold chilling by salicylic acid in banana seedlings. *Acta Bot Sin* 45: 567-573

Kang HM, Saltveit ME, 2001. Activity of enzymatic antioxidant defense systems in chilled and heat shocked cucumber seedling radicles. *Physiologia Plantarum* 113(4): 548-556

Kang HM, Saltveit ME, 2002. Chilling tolerance of maize, cucumber and rice seedling leaves and roots are differentially affected by salicylic acid. *Physiol Plant* 115: 571-576

Koch JR, Creelman RA, Eshita SM, Seskar M, Mullet JE, Davis KR, 2000. Ozone sensitivity in hybrid poplar correlates with insensitivity to both salicylic acid and jasmonic acid. The role of programmed cell death in lesion formation. *Plant Physiol* 123: 487-496

Kusumi K, Yaeno T, Kojo K, Hirayama M, Hirokawa D, Yara A, Iba K, 2006. The role of salicylic acid in the glutathione-mediated protection against photooxidative stress in rice. *Physiol Plant* 128: 651-666

Lederer B, Boger P, 2003. Antioxidative responses of tobacco expressing a bacterial glutathione reductase. *Z Naturforsch* 58c: 843-849

Leon J, Lawton MA, Raskin I, 1995. Hydrogen peroxide stimulates salicylic acid biosynthesis in tobacco. *Plant Physiol* 108: 1673-1678

Metwally A, Finkemeier I, Georgi M, Dietz KJ, 2003. Salicylic acid alleviates the cadmium toxicity in barley seedlings. *Plant Physiol* 132: 272-281

Mittler R, Feng XQ, Cohen M, 1998. Post-transcriptional suppression of cytosolic ascorbate peroxidase expression during pathogen-induced programmed cell death in tobacco. *Plant Cell* 10: 461-474

Molina A, Bueno P, Marin MC, Rodriguez-Rosales MP, Belver A, Venema K, Donaire JP, 2002. Involvement of endogenous salicylic acid content, lipoxygenase and antioxidant enzyme activities in the response of tomato cell suspension cultures to NaCl. *New Phytol* 156: 409-415

Munne-Bosch S, Penuelas J, 2003. Photo- and anti-oxidative protection, and a role for salicylic acid during drought and recovery in field-grown *Phillyrea angustifolia* plants. *Planta* 217: 758-766

Mutlu S, Atici Ö, 2013. Alleviation of high salt toxicity-induced oxidative damage by salicylic acid pre-treatment in two wheat cultivars. *Toxicol Industrial Health* 29(1): 89-96

Nawrath C, Metraux JP, 1999. Salicylic acid induction-deficient mutants of *Arabidopsis* express PR-2 and PR-5 and accumulate high levels of camalexin after pathogen inoculation. *Plant Cell* 11: 1393-1404

Nawrath C, Heck S, Parinthawong N, Metraux JP, 2002. EDS5, an essential component of salicylic acid-dependent signaling for disease resistance in *Arabidopsis*, is a member of the MATE transporter family. *Plant Cell* 14: 275-286

Nemeth M, Janda T, Horvath E, Paldi E, Szalai G, 2002. Exogenous salicylic acid increases polyamine content but may decrease drought tolerance in maize. *Plant Sci* 162: 569-574

Pal M, Horvath E, Janda T, Paldi E, Szalai G, 2005. Cadmium stimulates the accumulation of salicylic acid and its putative precursors in maize (*Zea mays*) plants. *Physiol Plant* 125: 356-364

Rao MV, Davis KR, 1999. Ozone-induced cell death occurs via two distinct mechanisms in *Arabidopsis*: the role of salicylic acid. *Plant J* 17: 603-614

Rao MV, Paliyath G, Ormrod DP, Murr DP, Watkins CB, 1997. Influence of salicylic acid on H₂O₂ production, oxidative stress, and H₂O₂-metabolizing enzymes. Salicylic acid-mediated oxidative damage requires H₂O₂. *Plant Physiol* 115(1): 137-149

Reddy KP, Subhani SM, Khan PA, Kumar KB, 1995. Effect of light and benzyl adenine on dark-treated growing rice leaves, II: Changes in peroxidase activity. *Plant Cell Physiol* 24: 987-994

Scandalios JG, 1997. Molecular genetics of superoxide dismutase in plants, pp. 527-568. In: *Oxidative Stress and the Molecular Biology of Antioxidant Defenses*. Scandalios JG ed. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press

Schmittgen TD, Livak K, 2008. Analyzing real-time PCR data by the comparative CT method. *Nature Protoc* 3(6): 1101-1108

Scott IM, Clarke SM, Wood JE, Mur LAJ, 2004. Salicylate accumulation inhibits growth at chilling temperature in *Arabidopsis*. *Plant Physiol* 135: 1040-1049

Shi Q, Bao Z, Zhu Z, Ying Q, Qian Q, 2006. Effects of different treatments of salicylic acid on heat tolerance, chlorophyll fluorescence, and antioxidant enzyme activity in seedlings of *Cucumis sativa* L. *Plant Growth Regul* 48: 127-135

Stevens J, Senaratna T, Sivasithamparam K, 2006. Salicylic acid induces salinity tolerance in tomato (*Lycopersicon esculentum* cv. Roma): associated changes in gas exchange, water relations and membrane stabilisation. *Plant Growth Regul* 49(1): 77-83

Strobel NE, Kuc JA, 1995. Chemical and biological inducers of systemic resistance to pathogens protect cucumber and tobacco plants from damage caused by paraquat and cupric chloride. *Phytopathol* 85: 1306-1310

Surplus SL, Jordan BR, Murphy AM, Carr JP, Thomas B, Mackerness SAH, 1998. Ultraviolet-B-induced responses in *Arabidopsis thaliana*: role of salicylic

acid and reactive oxygen species in the regulation of transcripts encoding photosynthetic and acidic pathogenesis-related proteins. *Plant Cell Environ* 21: 685–694

Tasgin E, Atici O, Nalbantoglu B, 2003. Effects of salicylic acid and cold on freezing tolerance in winter wheat leaves. *Plant Growth Regul* 41: 231–236

Van der Does, D, Leon-Reyes A, Koornneef A, Van Verk MC, Rodenburg N, Pauwels L, Pieterse, CM, 2013. Salicylic acid suppresses jasmonic acid signaling downstream of SCFCOI1-JAZ by targeting GCC promoter motifs via transcription factor ORA59. *Plant Cell* 25(2): 744-761

Vlot AC, Dempsey DA, Klessig DF, 2009. Salicylic acid, a multifaceted hormone to combat disease. *Annu Rev Phytopathol* 47: 177–206

Yadava, P, Kaur, P, Singh, I, 2013. Exogenous application of ascorbic acid alleviates oxidative stress in maize. *Indian J Plant Physiol* 18: 339-343

Yang YN, Qi M, Mei CS, 2004. Endogenous salicylic acid protects rice plants from oxidative damage caused by aging as well as biotic and abiotic stress. *Plant J* 40: 909–919

Yang ZM, Wang J, Wang SH, Xu LL, 2003. Salicylic acid-induced aluminum tolerance by modulation of citrate efflux from roots of *Cassia tora* L. *Planta* 217: 168–174

Yuan S, Lin HH, 2004. Transcription, translation, degradation and circadian clock. *Biochem Biophys Res Commun* 321: 1–6

Yusuf M, Hayat S, Alyemeni MN, Fariduddin Q, Ahmad A, 2013. Salicylic acid: Physiological roles in plants, pp. 15-30. In: Salicylic Acid. Springer Netherlands