

## Different mutations in the *ZmCAD2* gene underlie the maize brown-midrib1 (*bm1*) phenotype with similar effects on lignin characteristics and have potential interest for bioenergy production

Yves Barrière<sup>1\*</sup>, Hélène Chavignéau<sup>1</sup>, Sabrina Delaunay<sup>1</sup>, Audrey Courtial<sup>1,2,3</sup>, Mickael Bosio<sup>4</sup>, Hervé Lassagne<sup>4</sup>, Jérémy Derory<sup>4</sup>, Catherine Lapierre<sup>5</sup>, Valérie Méchin<sup>5</sup>, Christophe Tatout<sup>4\*</sup>

<sup>1</sup>INRA, Unité de Génétique et d'Amélioration des Plantes Fourragères, 86600 Lusignan, France

<sup>2</sup>Université de Toulouse, UPS, UMR5546, LRSV, BP 42617, Auzet, 31326 Castanet-Tolosan, France

<sup>3</sup>CNRS, UMR 5546, BP 42617, Auzet, 31326 Castanet-Tolosan, France

<sup>4</sup>Biogemma, site de la Garenne, CS 90126, 63720 Chappes, France

<sup>5</sup>INRA, Institut Jean-Pierre Bourgin, 78026 Versailles, France

\*Corresponding author: E-mail: yves.barriere@lusignan.inra.fr

<sup>#</sup>current address - CNRS UMR6293, INSERM U1103, GRED, les Cézeaux, 24 Avenue des Landais, BP 80026 Aubière cedex, France

### Abstract

The maize *ZmCAD2* gene has been fully sequenced in several normal and *bm1* maize lines, highlighting a large diversity of mutations underlying the *bm1* phenotype. Mutations in three *bm1* lines (F2bm1, A619bm1, and 511Jbm1) were found corresponding to short InDels inducing premature stop codons and truncated proteins. In two lines (511Kbm1 and 5803Cbm1), mutations were limited to an only SNP or to a few SNP, modifying the catalytic sites, and likely inactivating the proteins. Results also established that the 5803lbm7 mutant was in fact a *bm1* mutant, with a sequence fully identical to the 5803Cbm1 sequence. The two new F7803bm1 (natural mutant) and Ev2210bm1 (transposon tagging *Mutator* investigations) both had a transposon insertion in the *ZmCAD2* DNA, resulting in a truncated protein, even if the mRNA was produced. The biochemical characteristics of the Ev2210bm1 lignins corroborated the signature of CAD2 deficiency in plants, with the presence of aldehydes and atypical compounds and linkages. Considering lignin structure and content, CAD2 is likely a good target for the improvement of energy production based on maize and grass lignocellulosic biomass, including a greater susceptibility to environmentally friendly pretreatments, as it was shown in *bmr* sorghum. The interest in maize *bm1* hybrids for cattle feeding also should be considered because there seem to be little or limited negative effects of CAD2 mutations on other agronomical traits.

**Keywords:** maize, brown midrib, *bm1*, lignin, ferulic acid, cell wall, degradability, biofuel, biogas

### Introduction

Maize brown-midrib (*bm*) mutants exhibit a red-dish-brown pigmentation of the leaf midrib and stalk pith, associated with lignified tissues, after the plants have about five expanded leaves. The first *bm* mutants were described and investigated successively by Kiesselbach (1922), Eyster (1926), and Jorgenson (1931). The latter established that mutations of these first mutants were “due to identical factors” and that the “*bm* character” segregated as “a simple Mendelian recessive” trait. This gene was later named *bm1* when the *bm2* mutant was described by Burnham and Brinks (1932). A little later, two other genes inducing the *bm* phenotype were described as *bm3* (Emerson, 1935) and *bm4* (Burnham, 1947). The *bm2*, *bm3*, and *bm4* genes also originated from natural mutations and segregated as simple Mendelian recessive traits. The *bm1* mutation was assigned to maize chromosome 5 (bin 5.04) in position 77.5 cM, close to the centromere (Jorgenson, 1931; MaizeGDB, [www.maizegdb.org](http://www.maizegdb.org)). The three other *bm2*, *bm3*, and

*bm4* mutants were later mapped in bins 1.11, 4.05, and 9.07, respectively (MaizeGDB database). No new maize *bm* mutations were characterized for nearly 60 years after the Burnham’s paper (1947), despite the mention of additional natural *bm* mutants in the MaizeGDB database. Recent allelic tests of these latter mutants highlighted the three novel *bm5*, *bm6*, and *bm7* loci (Haney et al, 2008; Ali et al, 2010). While *bm5* and *bm7* mutations are still uncharacterized, the *bm6* mutations have been mapped to a 180 kb region of bin 2.01. Ten underlying candidates were proposed (Chen et al, 2012a), including a gene encoding a zinc finger CCCH-type protein, of which different family members also colocalized with cell wall degradability QTLs in RIL progenies of *Arabidopsis* (Chavignéau et al, 2012) and maize (Courtial et al, unpublished data).

In the final step of maize monolignol biosynthesis, the cinnamyl alcohol dehydrogenase (CAD) protein(s) catalyze(s) the reduction of *p*-coumaryl, coniferyl, and sinapyl aldehydes to their corresponding alcohols, using NADPH as a cofactor, prior to their trans-

port to the wall and their polymerization into the lignin polymer. Two types of CAD genes are currently considered in plants, based on original investigations in eucalyptus. EgCAD1-type enzymes are short-chain alcohol dehydrogenases (Jornvall et al, 1995; Goffner et al, 1998), which are active as monomers on coniferaldehyde, but not sinapaldehyde (Hawkins and Boudet 1994; Damiani et al, 2005). A ZmCAD1 activity has been described in maize (Kanazawa et al, 1999), corroborating the existence of the two types of CAD enzymes also in (this) grass species. EgCAD2-type enzymes are conversely zinc-containing long-chain alcohol dehydrogenases, which are active as dimers, and are considered the predominant CAD proteins involved in lignification (Jornvall et al, 1987; Goffner et al, 1992; Hawkins and Boudet, 1994). In maize *bm1* plants, the activity of a CAD enzyme was shown to be reduced by 60 - 70% in the above-ground organs and by 90 - 97% in roots (Halpin et al, 1998). These authors therefore considered that "*bm1* is not a null mutation of the *ZmCAD*, but affects its expression, possibly through alterations in upstream or downstream non-coding regions". Linkage analyses also showed that the *ZmCAD* locus was closely linked to RFLP markers corresponding to the position of the *bm1* mutation. Halpin et al (1998) thus concluded "that *ZmCAD* is very likely to be allelic to *bm1*". This maize *ZmCAD* gene associated with the *bm1* mutation was latter shown to be orthologous to *EgCAD2*, and it has thus been named *ZmCAD2* (Guillaumie et al, 2007b). However, a lack of complete specificity of their polyclonal antibody left the possibility of reactions with other CAD or CAD-like proteins, and/or only partial reactions against the targeted *ZmCAD2* protein. In addition, gene expression, which was investigated based on northern analysis with degenerate primers based on the tobacco CAD2 protein sequence, of which specificity against the *ZmCAD2* sequence is likely partial, indicated a residual CAD expression in *bm1* leaves and stems. In addition, based on gene expression investigations with the MaizeWall macroarray, the *ZmCAD2* gene was shown to be under-expressed in *bm1* plantlets with a residual expression of 0.36 as compared to the control F2 line (Guillaumie et al, 2007a). This agreed closely with the residual activity described by Halpin et al (1998). In addition, two *ZmCAD2*-like genes and the *ZmCAD1* gene were simultaneously under-expressed in F2bm1 plantlets. Similarly, based on Suppression Subtractive Hybridization (SSH) and micro-array data, Shi et al (2006) also established that several *ZmCAD* and CAD-like genes were down-regulated in young *bm1* plants (5-7 week-old plants).

The maize *ZmCAD2* gene (GRMZM5G844562) associated with the *bm1* mutation is located at position 98.993 Mbp on the chromosome 5 (bin 5.04) of the maize B73 genome (Schnable et al, 2009), upstream the estimated position of the chromosome 5 centromer. The maize *ZmCAD1* gene

(GRMZM2G179981) is also located in bin 5.04, in a downstream position (129.940 Mbp), upstream of the estimated position of the centromere. The latter gene only has distant paralogs, and at least one of them encodes a dihydroflavonol-4-reductase. The *ZmCAD2* also has no very close paralogs, and several of these *ZmCAD2*-like genes are now annotated as encoding mannitol dehydrogenase enzymes (maizesequence database, [www.maizesequence.org](http://www.maizesequence.org)). As a tentative conclusion, *ZmCAD1* and *ZmCAD2* are likely the two *ZmCAD* genes primarily involved in normal constitutive lignification of maize.

The consequences of the *bm1* mutation on maize lignin content and structure were first described forty years after the description of the mutation (Kuc and Nelson, 1964; Gee et al, 1968). Mature maize *bm1* plants have a lignin content that is reduced by 10 to 20%, a slight decrease in ferulic acid (FA) esters and substantially reduced contents (about 40%) in p-coumaric (pCA) esters and FA ethers (Provan et al, 1997; Barrière et al, 2004a). The frequency of p-hydroxyphenyl (H), guaiacyl (G) and syringyl (S) thio-acidolysis monomers was similar in *bm1* and normal plants, showing that the *bm1* mutation does not specifically affect one of the lignin units. However, the reduced recovery of thioacidolysis monomers reveals that the frequency of lignin units involved only in  $\beta$ -O-4 bonds was about 50% lower in *bm1* plants than in lignins of normal plants, indicating that lignins of *bm1* plants were substantially enriched in carbon-carbon inter-unit linkages (Halpin et al, 1998; Barrière et al, 2004a). Lignins of *bm1* plants are also typified by a substantial incorporation of coniferaldehyde and, to a lower extent, of sinapaldehyde and p-hydroxybenzaldehyde into the polymer (Jacquet, 1997; Kim et al, 2002; Kim et al, 2003; Barrière et al, 2004a). This notable incorporation of p-hydroxy-cinnamaldehyde-derived compounds in *bm1* lignins was in agreement with the under-expression of the *ZmCAD2* gene.

Concerns over global climate changes, together with a growing worldwide demand for energy, have highlighted the crucial need for alternative resources to replace fossil fuels. Second-generation biofuels, based on lignocellulose materials, have opened up new avenues, including a large valorization of agricultural and woody residues. The latter do not compete with food supplies, unlike biofuels based on cereal or oleaginous grains. The European Union commissioner for "Climate Action" Connie Hedegaard indeed recently said that "we cannot morally afford to build a very big industry on something that is not good for the environment or for food prices". However, the biological conversion of cell wall carbohydrates, mainly located in the secondary lignified plant cell walls, into fermentable sugars is hindered by their association with lignins, as well as with p-hydroxycinnamic acids in grasses. A similar situation has existed for decades in breeding programs towards higher digestibility and energy value of forage plants. Improvement of maize

energy value, for both cattle nutrition and industrial purposes, can be based on crosses and breeding with lines for which high values of cell wall degradability have been shown (Barrière et al, 2009; Barrière et al, 2010). However, the use of mutants with similarly higher cell wall degradability is also a relevant strategy in plant breeding. A mutant gene with major effect can indeed be more easily and rapidly back-crossed in elite lines based on marker assisted selection than several quantitative traits originating from one or more genetic resources.

Based on the B73 *ZmCAD2* sequence available from the Maize Genomics Sequencing Project (MGSP, [www.maizesequence.org](http://www.maizesequence.org)), the objective of this work was to sequence the *ZmCAD2* gene in *bm1* lines in comparison with their normal counterpart, including the reference line F2bm1, several new *bm1* mutants (Ali et al, 2010), and a new natural *bm1* mutant identified in an INRA nursery. In order to strengthen the involvement of the *ZmCAD2* gene in the *bm1* phenotype, a novel maize mutant obtained from transposon tagging in the *ZmCAD2* gene was simultaneously considered and characterized during these investigations. Transcriptomic approaches were also performed in order to estimate the respective effects of different *bm1* mutations on *ZmCAD* gene expression. In addition, because conflicting results were observed in new allelism tests, the *bm7* mutant was added to the sequencing investigations. Finally, the potential of the *bm1* mutation was discussed for further improvement of plant energy value for both cattle feeding and biofuel production.

## Materials and Methods

### Origins of *bm1* and *ZmCAD2* mutant lines

The F2bm1 line was obtained after seven back-crosses by the early flint INRA F2 line of the original *bm1* source (95033-47) supplied to INRA Lusignan by LF Bauman of Purdue University (West Lafayette, Indiana, USA) in 1972. This INRA F2bm1 genetic resource was also used in investigations by Halpin et al. (1998). Seeds of A619 and A619bm1 were supplied by N de Leon of University of Wisconsin (Madison) to INRA Lusignan in 2009. Seeds of the recently identified new *bm1* mutants 5803Cbm1, 511Kbm1 (or 5803Lbm1) and 511K, 511Jbm1 (or 5803Jbm1) were obtained in 2010 and 2011 from the Maize Genetics Cooperation Stock Center (MGCSC, [www.maizegdb.org](http://www.maizegdb.org)). The *ZmCAD2* mutant was isolated by the Biogemma company from transposon tagging investigations with the *Mutator* element, in the framework of the French national Génoplante program. Seeds of this mutant (*Zmcad2-m2210::Mu* named thereafter Ev2210bm1), and the corresponding normal control seeds (Ev2210), resulted from three generations of self-pollinations after five generations of backcrossing (BC5S3) the original mutant plant with an elite line of the Limagrain company. The new INRA F7803bm was found in 2009 in an S7 generation of a flint prog-

eny (Laborde J and Bauland C, pers comm). During the allelism tests of the F7803bm mutation, brown-midrib plants were unexpectedly obtained in both F7803bm x F2bm1 and F7803 x 5803Lbm7 crosses. Consequently, the 5803Lbm7 line was added to the current investigations.

### Biochemical characterization of the *ZmCAD2*

#### *Ev2210bm1* mutant

Plants of *Ev2210bm1* *ZmCAD2* mutant and their normal *Ev2210* isogenic were grown in 2007 summer nursery in Mons (Puy de Dôme, France) according to current maize cropping conditions. Cell wall and lignin analyses were carried out on whole plants without ears, harvested at silage stage maturity. Data were obtained from samples of nine normal and nine mutant plants (without ears). Two replicates were analyzed. Neutral detergent fiber (NDF, or extract-free samples) and acid detergent lignin (ADL) were measured according to Goering and Van Soest (1970). Klason lignin (KL) was estimated according to Dence and Lin (1992). KL values are usually two to four times greater in grasses than ADL estimates, with the loss of an acid-soluble lignin part in the first step of the ADL procedure (Hatfield et al, 1994; Jung et al, 1997; Hatfield and Fukushima, 2005). ADL and KL, which are constituents of the cell wall, were expressed as percentage of NDF (ADL/NDF and KL/NDF). The *in vitro* dry matter digestibility (IVDMD) was estimated according to Aufrère and Michalet-Doreau (1983). The cell wall digestibility was then estimated according to Struik (1983) and Dolstra and Medema (1990) as the *in vitro* NDF digestibility (IVNDFD) assuming that the non-NDF part of plant material was completely digestible [IVNDFD = 100 x (IVDMD - (100 - NDF)) / NDF]. Thioacidolysis was performed from extract-free samples and according to Mir Derikvand et al (2008). In addition, some thioacidolysis experiments were conducted on exhaustively permethylated samples in order to evaluate the proportion of terminal units with free phenolic groups relative to internal units, according to a previously described procedure (Lapiere et al, 1988).

### Allele sequencing and PCR analyses

#### DNA extraction and sequencing

DNA of each genotype was extracted from 1 g of fresh leaf using a modified CTAB protocol (CIM-MYT, 2005). Sequencing was performed for each PCR fragment in both directions by Millegen ([www.millegen.com](http://www.millegen.com), France). Contigs were constructed using Staden software [<http://staden.sourceforge.net/> (version v4.8b1)]. Sequences were aligned using MultAlin software [<http://multalin.toulouse.inra.fr/multalin/multalin.html> (Corpet, 1988)]. The deduced amino acid sequences were obtained using Transeq software (<http://mobyle.pasteur.fr/cgi-bin/portal.py?#forms::transeq>). Amino acid sequences were aligned similarly using MultAlin software.

**Table 1** - Primer pairs designed for amplification of the whole DNA *ZmCAD2* gene, and for expression studies, based on the B73 DNA sequence.

Forward primer (5'-> 3')	Reverse primer (5'-> 3')	Length (bp)
<b>Gene sequencing</b>		
ZmCAD2_1F1 GACCTCCTCGAAAGACGAAA	ZmCAD2_1R1 TGTCCCATCGAAATGTTGAA	1,256
ZmCAD2_2F1 TAATTGGGGGAGCACTCTG	ZmCAD2_2R1 ATAGCTCCGGACTGGGATG	1,281
ZmCAD2_3F1 CTCCGTGTACGGTCAGAAT	ZmCAD2_3R1 TGGTGGTTAACCTCACAAA	1,279
ZmCAD2_3F2 CATGACGGACAGGACAAACAC	ZmCAD2_3R2 GCATGCAAGATAACGCTGAA	446
ZmCAD2_4F1 CTTATGGTGACCAGGCAAGAG	ZmCAD2_4R1 GACGCAGAACTGAAGCACCT	999
ZmCAD2_5F1 CGACTCGCTGGACTACATCA	ZmCAD2_5R1 GGTGCGGGTATTGCTCAAG	967
<b>Expression studies</b>		
ZmCAD1aF CTTCCCTTCCGGACAAGTGTG	ZmCAD1aR AGCTTTCGATGGTCTCCTTGACG	-
ZmCAD2cF GGTACATACAACGACGTCTACACTG	ZmCAD2cR CCGGGATCTTCACCACAAACTTC	-
ZmCAD4bF TCAAGCCAACGGCAAGATG	ZmCAD4bR CAGGGTCTGTTCCCAATGATGAG	-
ZmUbiquitinF CATTGTGCCCTGTTGAAC	ZmUbiquitinR AACAGCAACACCACAAACCA	-

#### Primer design and PCR amplifications

Pairs of primers used for the *ZmCAD2* gene sequencing were designed based on the available maize B73 sequence (GRMZM5G844562, [www.maizesequence.org](http://www.maizesequence.org), Table 1). Fragments of nearly 0.4 to 1.5 kb were amplified, encompassing the 5'UTR, the complete coding region, 1,000 bp before start codon and 1,000 bp after stop codon, giving a *ZmCAD2* investigated region close to 5 kb long. PCR reactions were performed in a final volume of 50 µl containing 1X PCR buffer and 0.5 U Taq DNA Pol (both MP Biomedicals, ref EPTQX925), 200 µM of dNTPs, 0.32 µM of 5' oligo, and 0.32 µM of 3' oligo. Fifty ng of genomic DNA was used as template. The program comprised 4 min at 94°C, followed by 35 cycles of 30 s at 95°C, 1 min at 60°C, and 1 min at 72°C, followed by 6 min at 72°C.

#### Expression analyses

For all eight normal (F2, 511K, Ev2210) and bm1 (F2bm1, 511Kbm1, 5803Cbm1, 511Jbm1, Ev2210bm1) genotypes, plants were grown in a glasshouse at INRA Lusignan in spring 2011. Lignin pathway genes had previously been shown to be highly expressed in the below-ear internode from tassel emergence to a few days after silking (Riboulet et al, 2009), a period corresponding to the end of the elongation phase and the deposition of the secondary cell wall in this internode. Consequently, below-ear internodes (without nodes) of three representative plants were harvested when tassels were clearly emerging out of the whorl. Samples were collected five hours after sunrise. To limit experiment costs, all plants of each line were pooled after cutting internodes into fragments nearly 1 cm long. All samples were immediately frozen in liquid nitrogen and stored at -80°C.

Total RNA was extracted and treated using the plant RNeasy mini kit and RNase-Free DNase Set (Qiagen, ref 70903 and 79254). RNA (1 µg) was reverse transcribed with 200 U of SuperScript III using 5µM oligo(dT) (Invitrogen, SuperScript III First-Strand

Synthesis System for RT-PCR, ref. 18080-051) according to the manufacturer's instructions. The resulting cDNAs was diluted 1:10. In addition to the *ZmUbiquitin* (GRMZM2G102471) gene used as standard, expression was investigated for the two *ZmCAD1* and *ZmCAD2* genes, and also for the *ZmCAD4* gene (GRMZM2G118610), which was originally annotated as a *ZmCAD2*-like gene (Barrière et al, 2009), but which is now considered to encode a mannitol dehydrogenase. Gene primer pairs were designed using the QuantPrime software to anneal near the 3' end of each transcript, often in 3'UTR area, ensuring primer specificity (Table 1). Real-time reverse transcription RT-PCR was performed on a LightCycler® (Roche Diagnostics) using the SYBR Green I Master (Roche Diagnostics, ref 04707516001). The PCR reaction mixture (20 µl) consisted of 3 µl water, 10 µl SYBR Green I Master Mix, 250 nM of each primer and 5 µl diluted cDNA template. The following LightCycler experimental run protocol was used including a denaturation program (95°C for 10 min), an amplification and quantification program repeated 45 times (95°C for 10 s, 60°C for 15 s, 72°C for 25 s with a single fluorescence measurement), a melting curve program (65-95°C with a heating rate of 2.2°C per second and a continuous fluorescence measurement), and finally a cooling step to 40°C. The PCR threshold cycle number of each gene was normalized with that of the *ZmUbiquitin* reference gene to calculate the relative mRNA levels. The Fisher test (P < 0.1) was used for statistical analyses of quantitative RT-PCR data.

## Results

### Allele sequencing of the *ZmCAD2* gene in normal and bm1near-isogenic lines

#### F2 and F2bm1 lines

The F2 *ZmCAD2* allele differed from the B73 *ZmCAD2* allele in the 3' region, with 12 SNPs and two 54- and 30-bp long deletions starting from positions 4,031 and 4,116 (relative to the transcriptional start site in B73), respectively. In addition, amino

acid 349 is an asparagine in F2, while it is an aspartic acid in B73. The *ZmCAD2* alleles in B73 and F2bm1 were overall similar, likely related to the fact that the original *bm1* mutation used in F2 backcrossing occurred in a dent Corn Belt line and not in a European flint one. However, a two basepair (AC) insertion was identified in exon 3 of the F2bm1 line after the C base located in B73 CDS position 392 (Supplementary Figure 1). The ORF was consequently disrupted in F2bm1 by a premature stop codon, resulting in a predicted truncated protein of 147 amino acids, in comparison to the 367 amino acid long proteins in B73 and F2 (Supplementary Figure 2). Such a truncated *ZmCAD2* protein, in which the site involved in enzyme binding to its substrate, a catalytic site, and several binding sites to cofactors, were missing is very likely to be non functional, even if produced. The AC base insertion may therefore explain the *bm1* phenotype of the F2bm1 line. In addition, in F2, F2bm1, as in all investigated lines, in B73 CDS position 951, there was a synonymous substitution of an A base (B73) by a G base (all lines), the GGA and GGG codons both encoding for a glycine.

#### *A619 and A619bm1 lines*

Allele sequencing was only investigated in the area overlapping the AC insertion shown in F2bm1. The same mutation was highlighted in A619bm1 as the one identified in F2bm1. This suggests that the backcrosses of the *bm1* mutation in the A619 line were made with the same *bm1* mutant allele as the one used for the F2bm1 line, which is, given the origin of these lines, likely the original *bm1*-reference (*bm1-ref*) allele.

#### *511Jbm1 line*

A CGCG four base insertion was identified in the *ZmCAD2* exon 3 of the 511Jbm1 line, which occurred after the CG bases located in positions 366-367 of the B73 CDS (Supplementary Figure 1, *bm1-J* allele). As observed in F2bm1, this insertion disrupted the ORF and induced a 127 amino acid long truncated protein (Supplementary Figure 2), which is very likely non functional, even if produced.

#### *511Kbm1 line*

Allelic variation in the *ZmCAD2* CDS of the 511Kbm1 line, in comparison to B73, was reduced to only one SNP in exon 3 substituting the GGA codon into a GAA codon (CDS bp 401), thus substituting a glycine (G) by a glutamic acid (E) as the 134<sup>th</sup> amino acid of the *ZmCAD2* protein (Supplementary Figures 1 and 2, *bm1-K* allele). This unique mutation occurred in an area encoding the between-species conserved <sup>130</sup>PTQGGFA<sup>136</sup> amino acid motif (<sup>130</sup>PTQGEFA<sup>136</sup> in the mutant), located just before the  $\beta$ 9 strand. However, a 21 bp deletion was also shown in the 5'UTR, in position 143-123 bp before the B73 ATG start codon, in which was inserted a 5 bp substitution establishing an extra TATATA box 128 bp before the *ZmCAD2* ATG start codon of the 511K line, in addition to the TATATA box located 40 bp upstream. These two


events were shown to be specific to the 511Kbm1 line and were not present in the 511K near-isogenic normal line. It was thus considered that either changes in protein structures inactivated its catalytic activity, or less likely that the transcription was reduced to a quasi-null level due to the extra TATA box.

#### *5803Cbm1 line*

Neither insertions nor deletions were shown in the *ZmCAD2* CDS of the 5803Cbm1 line, in comparison to B73, but the two sequences differed by two synonymous and three non synonymous SNPs. Synonymous SNPs occurred in CDS positions 192 bp (CCT/CCC, proline) and 237 bp (GGG/GGA, glycine). The first non-synonymous SNP was shown in exon 1 where a TCC codon was substituted by an ACC codon (CDS bp 76), encoding a threonine (T) instead of a serine (S) as the *ZmCAD2* 26th amino acid (Supplementary Figures 1 and 2). The second non-synonymous SNP occurred in exon 3, where the TCC codon was changed into TGC (CDS bp 410), encoding a cysteine (C) in place of a serine (S) as the 137th amino acid. The latter mutation, and the one shown in exon 1, likely did not greatly change activities of the encoded enzyme, as they both occurred in between-species genomic unconserved areas (Youn et al, 2006). The third mutation also occurred in exon 3, with a codon CAC changed into CGC (CDS bp 205, which substituted a histidine (H) by an arginine (R) as 69<sup>th</sup> amino acid in the conserved amino acid motif <sup>68</sup>GHEVVGXXXXGXXV<sup>82</sup> leading to a <sup>68</sup>GREVVG<sup>73</sup> motif in the mutant (Supplementary Figures 1 and 2, *bm1-C* allele). The latter motif is involved in the binding of the zinc ion at the catalytic site of the enzyme, and the GHE conserved amino acid sequence also corresponded to the area separating the  $\beta$ 4 and  $\beta$ 5 strands (Youn et al, 2006). This H to R mutation probably alters the catalytic activity of the encoded enzyme, and is consequently likely the cause of the *bm1* phenotype. In addition, a very important polymorphism was shown in the *ZmCAD2* line first intron of the 5803Cbm1 line, which differed from B73 first intron by 54 SNP and 18 Indels. These changes could also modify the *ZmCAD2* gene expression in 5803Cbm1 mutant plants. Finally, the 5'UTR of the *ZmCAD2* gene in 5803Cbm1 mutant line exhibited the same 15 bp deletion and extra TATATA box as observed in the 511Kbm1 mutant line.

#### *5803Ibm7 line*

The sequence of the *ZmCAD2* gene in the 5803Ibm7 line was found to be fully identical to the sequence of the 5803Cbm1 line. This result was in agreement with new allelism tests done at INRA Lutignan showing that the 5803Ibm7 line gave hybrids with brown midribs in crosses with F2bm1 (Figure 1). Allelism between *bm1* and *bm7* was corroborated by another allelism test done by Hongjun Liu (Iowa State University, [pers com](#)) and it was also in agreement with unpublished previous observations of Sarah Hake (2007, [pers com](#)). The mutation in the 5803Ibm7



**Figure 1** - Allelism tests between 5807bm7 and F2bm1 (A), F7803bm and F2bm1 (B), F7803bm and 5803bm7 (C), Ev2210bm and F2bm1 (E, F), all four crosses giving brown-midribs, and F7803bm and F2bm3 (D), giving normal midribs.

line is definitely allelic to *bm1* (*bm1-7* allele) and this fact is now specified in the maizeGDB database ([www.maizeGDB.org](http://www.maizegdb.org), locus *bm7*).

#### *F7803bm1* line

Brown-midrib plants were obtained in crosses of F7803bm with F2bm1 (Figure 1), but not in crosses with F2bm2, F2bm3, or F2bm4, and similarly not with the newly described *bm5* and *bm6* lines. However, the F7803bm line also gave plants with brown

midribs in crosses with 5803bm7 (Figure 1). This fact further validated the allelism of the two *bm1* and *bm7* mutations. In contrast to investigations with the five previous *bm1* mutants, the sequence of the *ZmCAD2* gene in the F7803bm1 line was not fully obtained, with a failure in amplifying the whole second exon. PCR products corresponding to the second exon were small, with *ZmCAD2* homologies reduced to a few base pairs, which most often cor-

responded to primer sequence. The forward primer used was located 459 bp upstream of exon 2, and the reverse primer was located 629 bp downstream of exon 2 (these two primers enabled amplification of exon 2 in normal lines). The most probable situation in F7803bm1 would therefore be an insertion of a (retro-) transposon element in the exon 2 area, preventing PCR. In addition, there was an amino acid substitution in exon 4, with a glutamic acid replaced by an aspartic acid, corresponding in B73 CDS position 924 to a substitution of a G nucleotide by a T nucleotide (Supplementary Figures 1 and 2, allele *bm1-F*). Glutamic and aspartic acids are both diacid amino acids, and such a change is not expected to be responsible for the *bm1* phenotype. Nevertheless, the probable insertion of a (retro-) transposon close to exon 2 would very likely result in a protein disrupted before the region encoded by exon 4.

#### Ev2210bm1 line

In contrast to previous spontaneously occurring *bm1* mutations, the Ev2210 *ZmCAD2* *bm* mutant line was obtained from transposon tagging investigations with the Mutator element. Allelism tests showed that brown-midrib plants were obtained in crosses with F2bm1 (Figure 1), but not in crosses with F2bm2, F2bm3, or F2bm4, corroborating the involvement of *ZmCAD2* mutations in *bm1* phenotype. As it was the case for F7803bm1, the full sequence of the *ZmCAD2* gene in the Ev2210bm1 line was also not obtained. However, 24 bp of the upstream part and 26 bp of the downstream part of the *Mutator* element sequences


were obtained, showing that the *Mutator* element was located in this allele in exon 4, between position 678 and 679 bp of the B73 CDS. Lengths of a single copy of *Mutator* transposons are nearly 1,500 bp long, and they have a 220 bp long conserved terminally inverted repeats (TIRs). Such a long insertion, and chiefly the presence of the TIRs which cause very stable hairpin structures, would prevent an efficient PCR running. In any case, the *Mutator* element insertion induced a stop codon and disrupted the ORF, resulting in a 237 amino acid long truncated protein (Supplementary Figures 1 and 2, allele *bm1-M*), which is likely to be inactive if it is produced.

#### Gene expression in *bm1* mutants and their normal counterparts

The *ZmCAD2* gene appeared to be significantly under-expressed in all investigated *bm1* lines (F2bm1, 511Kbm1, 5803Cbm1, 511Jbm1, Ev2210bm1), in comparison to their normal counterparts (F2, 511K, Ev2210), corroborating the effect of *bm1* mutations on the expression of the *ZmCAD2* gene. In addition, the expression of the *ZmCAD1* gene was significantly up-regulated in three *bm1* mutants out of five, and especially in the Ev2210bm1 mutant line (Figure 2). These results did not corroborate the lower *ZmCAD1* expression observed in *bm1* plants (Shi et al., 2006; Guillaumie et al., 2007a), that was, however, observed in much younger plants. The *ZmCAD4* gene, encoding a CAD-like protein with a mannitol dehydrogenase activity, had variable expression levels according to genotypes. *ZmCAD4* was over-expressed in F2bm1 and Ev2210bm1 lines, while it was under-expressed in the three other *bm1* lines considered.

#### Lignin content and structure in the *ZmCAD2* Ev2210bm1 mutant line

The consequences of the *bm1* mutation on maize lignin content and structure have been shown to date in F2bm1 (*bm1-ref*) mutant plants (with the AC insertion), before it was firmly established that the *ZmCAD2* gene was affected in these plants. Relative to the wild-type plants, investigations in the Ev2210bm1 mutant plants, altered in the *ZmCAD2*



**Figure 2** - Comparative expression of *ZmCAD2*, *ZmCAD1*, and *ZmCAD4* genes in normal and *bm1* maize plants (\* =  $P < 0.1$ ; \*\* =  $P < 0.01$ ; \*\*\* =  $P < 0.001$ , and ns = non significant differences). The *ZmUbiquitin* gene was used as standard.

**Table 2** - Cell wall traits in whole plants without ear at silage maturity stage for the normal and *ZmCAD2* Ev2210bm1 mutant plants.

	NDF	KI/NDF	ADL/NDF	IVNDFD
Normal	61.15	15.64	6.14	34.59
Ev2210bm1	62.01	14.55	5.24	39.27
F test	3.0 ns	15.6 *	53.4 **	26.0 **

NDF (neutral detergent fiber) as % dry matter (DM), ADL (acid detergent lignin) and Kason lignin (KI) % NDF, *in vitro* NDF digestibility (IVNDFD) as weight % of NDF. Data as mean values from 9 independent samples, differences between normal and mutant plants were non significant (ns) or significant at  $P < 0.01$  % (\*) and  $0.001$  % (\*\*).

gene after transposon insertion, established that this mutant also displayed substantial variations in lignins and cell wall phenolics (Table 2). The lignin contents estimated by contents the Klason procedure and Van Soest (ADL) method were reduced by 6 and 17%, respectively, in Ev2210bm1 plants, while the cell wall content (referred to as NDF) was not changed. Substantial changes in lignin structure were also observed, with 30 - 40% fewer units only involved in  $\beta$ -O-4 bonds, as shown by the lower thioacidolysis yield (Table 3). Lignins in Ev2210bm1 plants are enriched in resistant inter-unit bonds, referred to as condensed linkages. Based on thioacidolysis investigations, the monomeric composition of the polymer was only slightly modified with a small increase in S unit frequency, a small decrease in G unit frequency, and no changes in H unit frequency. In addition to lignin-derived monomers, thioacidolysis of extractive-free samples released pCA and FA acids (Table 3). Previous studies have revealed that pCA is essentially ester-linked to lignins in maize cell walls (Ralph et al, 1994) and that most of these ester bonds do not survive the thioacidolysis procedure (Jacquet, 1997). Accordingly, the yield of thioacidolysis-released pCA accurately reflects the amount of *p*-coumarate esters. The origin of thioacidolysis-released FA is more complex because ferulate esters, which are linked to arabinoxylans, may be simultaneously linked to lignins by various bonding patterns. Anyway, obtained results suggested that the level of *p*-coumarate esters was substantially reduced in the cell walls of the Ev2210bm1 plants (nearly 30%), whereas ferulate units released by thioacidolysis were affected to a lower extent. Correlatively to lignin content and structure changes in Ev2210bm1 plants, their cell wall digestibility was improved by nearly 12%.

Following thioacidolysis and gas chromatography-mass spectrometry analysis of the lignin-derived monomers, signatures of CAD deficiency were identified in the Ev2210bm1 plants as previously observed in various CAD-deficient transgenic or mutant plants (Kim et al, 2002; Sibout et al, 2005; Thévenin et al, 2011). The mutant samples released

2 to 3 times more thioacidolysis monomers derived from coniferaldehyde end-groups than the control (Table 3). Moreover, the two G and S indene derivatives derived from coniferaldehyde and sinapaldehyde, which have been incorporated into lignins by 8-O-4 cross-coupling, were recovered in substantial amounts from Ev2210bm1 lignins and only as trace components from the control. These results corroborated those obtained in F2bm1 plants (Barrière et al, 2004a), even if aldehyde incorporations were shown to be lower in Ev2210bm1 plants than previously observed in F2bm1 plants. The difference between F2bm1 and Ev2210bm1 plants was probably related either to different penetrances of the two different types of *ZmCAD2* mutations in F2bm1 and Ev2210bm1, and/or to the different genetic backgrounds of the two lines. Among the differences, the much higher *ZmCAD1* over-expression was observed in Ev2210bm1, but not in F2bm1, plants (Figure 2).

Finally, when subjected to thioacidolysis and relatively to normal plants, Ev2210bm1 plants released 2 to 3 times higher amounts of the AG compound, which is a marker for the incorporation of free ferulic acid into lignins (Ralph et al, 2008). Despite this fact was not considered in previous years (Barrière et al, 2004a), a greater incorporation also occurred in F2bm1 lignins (data not shown), as well as in lignins of *Arabidopsis CAD-C x CAD-D* double mutant plants (Thévenin et al, 2011). The incorporation of ferulic acid into lignins allows new branching points with bi- and di-phenyl ether structures and consequently modifies the macro-molecular organization of the polymer (Ralph et al, 2008). The putative higher degree of branching of lignins in the Ev2210bm1 mutant was confirmed by the results obtained by thioacidolysis of permethylated samples which provides the proportion of terminal units with free phenolic groups relative to internal  $\beta$ -O-4 linked lignin units. In agreement with previous results obtained with maize lignins (Lapierre, 1993), thioacidolysis of the permethylated normal maize sample revealed that about 50% of G units are terminal units with free phenolic groups (Table 3), whereas this figure is higher for H

**Table 3** - Impact of the *ZmCAD2* Ev2210bm1 mutation on cell wall phenolics in whole plant without ear at the silage maturity stage based on thioacidolysis investigations.

Line	Thioacidolysis yield (H+G+S) ( $\mu\text{mole g}^{-1} \text{KL}$ )	Relative frequency of thioacidolysis monomers (% molar)			Minor thioacidolysis monomers ( $\mu\text{mole g}^{-1} \text{KL}$ )				Free phenolic groups (%) in $\beta$ -O-4 linkage			Thioacidolysis release <i>p</i> -OH cinnamic acids ( $\mu\text{mole g}^{-1} \text{KL}$ )	
		H units	G units	S units	coniferald end-groups	G Indene	S Indene	A <sub>6</sub>	H	G	S	pCA	FA
Normal	526.22	1.62	46.14	52.21	0.36	traces	traces	0.42	86.6	51.9	6.6	581.67	226.44
Ev2210bm1	294.89	1.60	43.57	54.82	1.18	9.54	7.27	1.27	91.5	61.3	7.8	420.22	218.89
F test	103.8**	0.2ns	12.0*	12.0*	172.4**	-	-	195.1**	6.4ns	161.3**	28.2**	40.4**	0.5ns

Thioacidolysis released H, G and S lignin monomers of coniferaldehyde (coniferald) end-groups, coniferaldehyde ether-linked at C $\beta$  (releasing G indene), sinapaldehyde ether-linked at C $\beta$  (releasing S indene), and of the marker compound (A<sub>6</sub>) for free ferulic acid incorporation in lignins. Thioacidolysis-released *p*-coumaric acid (pCA) and ferulic acid (FA). Percentage (%) of free phenolic groups in  $\beta$ -O-4 linked H, G and S lignin units, as determined by thioacidolysis of permethylated samples. Significant differences between normal and mutant plants indicated as in Table 2.

units (about 87%) and much lower for S units (less than 7%). However, in the Ev2210bm1 mutant, more than 60% of G lignin units are terminal units with free phenolic groups. The substantially higher frequency of free phenolic groups in lignins is a structural specificity which is common to all the CAD-deficient plants analyzed so far, including the F2bm1 maize mutant (Lapierre, 2010), and the signature of a lignin network disorganized in smaller domains.

## Discussion

While the *ZmCOMT* mutations were identified about 15 years ago in the maize *bm3* mutants (Vignols et al, 1995; Morrow et al, 1997), no investigations had firmly established up until now what gene was affected in *bm1* plants, even if it was greatly suspected that the *CAD2* gene was involved in the mutation(s). The sequencing of the *ZmCAD2* allele in eight *bm1* lines highlighted the diversity of genomic events underlying the *bm1* phenotype. Two- and four-base insertions, missense mutations, and transposon insertions are expected to lead to truncated or inactive CAD2 proteins, thereby causing the *bm1* phenotype. The mutation of the *ZmCAD2* gene in *bm1* maize plants was also established in concomitant investigations in other genetic backgrounds (Chen et al, 2012b). The first intron of the *ZmCAD2* *bm1-das1* mutant (Dow AgroSciences *bm1*) was shown to contain a 3,444 bp insertion, resulting in a chimeric mRNA containing a premature stop codon, and a truncated protein of 48 amino acids. The mutation identified in the *bm1* allele of the 515Dbm1 simultaneously investigated line was the same as the one observed in the F2bm1 and A619bm1 lines, consisting in an AC insertion in exon 3 that resulted in a 147 amino acid reduced protein. The AC insertion in the third *ZmCAD2* exon could thus be considered to be the first described *bm1* (*bm1-ref*) mutation.

In addition, based on observations in nurseries and allelism tests, the *CAD2-bm1* mutation seems to occur or appear more frequently than other maize *bm* mutations. This fact considered from a very small number of events could actually correspond to a random effect, without any underlying biological mechanism. However, this fact could also be related i) to the complexity of the catalytic sites of the CAD2 protein rapidly altered by a slight modification of the amino acid sequence, ii) possibly to a DNA sequence or a chromosomal position favoring transposon insertion, and iii) to a low or null effect of the *CAD2* mutation on agronomic value. Chen et al (2012b) have indeed observed that the CAG to TAG nonsense mutation occurred in *SbCAD2* of *bmr6* sorghum just one nucleotide apart from the AC insertion shown in maize *bm1* mutants. Moreover, although plants contain numerous genes coding for CAD-like genes, only *EgCAD2*-type, and to a lesser extent *EgCAD1*-type CAD enzymes, were shown with a primary physiological role in lignin biosynthesis. This situation results

from the presence in lignin-related CAD proteins of a few key residues permitting significant catalytic rates on monolignol precursors. The role of several of these key residues for efficient catalysis and monolignol biosynthesis had been especially established based on investigations in lead tree (*Leucaena leucocephala*) and switchgrass (Pandey et al, 2011; Saathoff et al, 2012). Correlatively, while a highly truncated protein was shown to be the determinant of the *SbCAD2* *bmr6-ref* mutation in sorghum, the *bmr6-3* mutation indeed resulted from a unique mutation in the <sup>188</sup>GXGG(V/L)<sup>193</sup> motif changed into a <sup>188</sup>GXGS(V/L)<sup>193</sup> motif, affecting the binding affinity for the NADP+ cofactor (Saballos et al, 2009). In addition, the rice *gold-hull-and-internode-2* (*gh2*) mutant, which exhibits a reddish-brown pigmentation in the hull and internodes which become golden yellow at maturation, has an altered *OsCAD2* gene with a G to A substitution in the fourth exon (Zhang et al, 2006). This substitution induced the replacement of a zero-charge glycine by an electro-negative aspartic acid, leading to an inactive protein, as confirmed by recombinant protein experiments. As observed in maize and sorghum, the CAD activity was drastically reduced in rice *gh2* mutant, indicating that other CAD(-like) enzymes only partly substitute the *OsCAD2* deficiency, at least in rice for coniferyl alcohol biosynthesis (Zhang et al, 2006).

All cell wall traits shown in plants of the Ev2210bm1 mutant were indeed very similar to the ones observed in the F2bm1 plants (Barrière et al, 2004a), corroborating the involvement of *ZmCAD2* mutations in the *bm1* phenotype. Both F2bm1 and the Ev2210bm1 plants had reduced contents in lignins and in *p*-coumarate esters, which are correlated with a higher cell wall digestibility. Both exhibited important structural alterations of their lignins, which correspond to the specific signature of CAD deficiency, namely a higher frequency of condensed bonds and of free phenolic groups and the increased incorporation of coniferaldehyde, sinapaldehyde and free ferulic acid in lignins (Lapierre et al, 2004; Ralph et al, 2008). However, in addition to the incorporation of aldehydes and atypical compounds into lignins, thioacidolysis yields indicated the synthesis of regular monolignols in *CAD2* mutant plants, reaching nearly 55 % of the normal  $\beta$ -O-4-linked monomeric units in the Ev2210bm1 maize mutant. *EgCAD1*-type and CAD-like genes acting on *p*-hydroxy-cinnamaldehydes, in addition to a possible residual CAD2 activity in mutant with modified but not truncated protein, should therefore be considered. Expression data showed a probable substitution role of the *ZmCAD1* gene in Ev2210bm1 and, to a lesser extent, in F2bm1 and 511Kbm1. Corroborating the substitution by other CAD or CAD-like enzymes, the truncated *SbCAD2* protein was not detected in extracts from *bmr6* sorghum plants, but CAD activity was still detectable in mutant plant tissues (Pillonel et al, 1991; Saballos et al, 2009). Nev-

ertheless, if other CAD proteins (including *EgCAD1*-type) can be active on cinnamyl substrates, brown midrib or golden phenotypes and deep alterations in lignin composition establish that *EgCAD2*-type genes encode the main CAD proteins involved in the regular monolignol biosynthetic pathway of grasses.

The interest of CAD2-deficient plants in industrial processes was first shown with the demonstration of the significantly improved pulping capacity of poplar and pine *CAD* mutants (Baucher et al, 1996; Lapierre et al, 1999; Lapierre et al, 2000; O'Connell et al, 2002; Gill et al, 2003). In grasses, for bioenergy production, the efficiency of the *EgCAD2*-type mutation or deregulation towards significant increases of enzymatic hydrolysis and/or conversion rate into bioethanol (with or without pretreatment) has been shown in sorghum and switchgrass plants, without significant negative effects on agronomic value (Saballos et al, 2008; Corredor et al, 2009; Sattler et al, 2009; Sattler et al, 2010a; Sattler et al, 2010b; Scott et al, 2010; Saathoff et al, 2011). In maize, most investigations with *bm* genes have been done with the *bm3* *COMT* mutant, because this latter induced the highest improvement in cell wall digestibility and energy value for cattle feeding (Barrière et al, 2004b). Similarly, high increases in the release of fermentable sugars were shown in *bm3* mutant plants (Vermeris et al, 2007). However, an important decrease in agronomic value has been simultaneously shown in nearly all early *bm3* investigated hybrids, a fact that could possibly be closely related to the germplasm of these ancient investigated hybrids. None of the considered hybrids were related to the lodent genetic resource, of high agronomic value and standability. Several *bm3* hybrids are now indeed available in different markets, including the US market, with proven greater efficiency in dairy cow feeding (at least 28 published investigations between 1976 and 2012, most of them in *Journal of Dairy Science*). The feeding value of maize *bm1* hybrids has not been extensively investigated, likely as the first digestibility measurements have shown the lower improvement of maize *bm1* genotypes in comparison to *bm3* near-isogenics (Barnes et al, 1971; Lechtenberg et al, 1972; Barrière and Argillier, 1993). Nevertheless, the agronomic value of maize *bm1* hybrids (and *CAD2* mutants) is seemingly less modified than the one of *bm3* hybrids (and *COMT* mutants). Based on INRA Lusignan unpublished data from the two old INRA260 and LG11 hybrids, average *in vivo* NDF digestibility values were equal to 50.6, 55.7, and 59.4% in normal, *bm1*, and *bm3* isogenics, while DM yields were equal to 12.7, 12.5, and 11.8 t ha<sup>-1</sup>, respectively. On the contrary, recent investigations strengthened the interest of maize *bm1*-type mutants, based on CAD down-regulation, for industrial purposes. Transgenic CAD-RNAi maize plants had stem cell walls with a slight reduction of lignin content, but the stems of deregulated plants were nevertheless more degradable than their

normal counterparts. Fermentation assays also revealed that CAD deregulated plants produced higher levels of ethanol compared to normal ones (Fornalé et al, 2012). Moreover, the degree to which lignin polymers incorporate various phenolic compounds in place of the three regular constitutive monolignols is surely underappreciated (Ralph, 2010). In addition to ferulic acid and hydroxy-cinnamaldehydes, unusual monomers including acylated hydroxycinnamyl alcohols, dihydro-hydroxycinnamyl alcohols, hydroxybenzaldehydes and other hydroxycinnamic acids, can be incorporated into lignins of wild-type plants (Vanholme et al, 2012). Because plants could tolerate shifts in lignin composition with no or lower impact on growth than observed for reduced lignin contents, the substitution of some fraction of the three regular monolignols by unusual or alternative monomers through breeding, directed mutagenesis, or genetic engineering is thus a relevant strategy to tailor lignins in bioenergy crops so that cell walls would be more susceptible to biomass pretreatments (Eudes et al, 2012; Vanholme et al, 2012). The interest of unusual monomers incorporation into lignins for bioenergy production was also highlighted with *bmr6* sorghum. Glucose yields were indeed improved by at least 25% from sorghum biomass of *bmr6* plants, compared to normal isogenics (Saballos et al, 2008; Dien et al, 2011). Several *bmr6* (and "bmr") sorghum hybrids have thus been registered, or are in registration process, for both European and American markets. The higher efficiency of *bmr6* sorghum for animal feeding has also been shown from dairy cows experiments. Cows fed *bmr6* silage in their diets had 16% higher milk yields in comparison to similar diet with normal sorghum (Oliver et al, 2003). However, milk yields were equal for *bmr6* sorghum and normal maize silages, highlighting again the higher cell wall degradability in maize than in sorghum. In addition, the modification of the lignin structure could also be considered to increase cell wall degradability (Zhang et al, 2011). In a set of maize inbred lines, more condensed lignins were thus shown to be more favorable to an increased cell wall degradability than  $\beta$ -O-4 rich lignins. Finally, for both bioenergy production and cattle feeding, the successful breeding of improved maize (and C4 grasses) genotypes with more degradable cell walls, based on different lignin polymer organization, incorporation of alternative monomers, and reduced ferulate cross-linkages, requires more knowledge about phenolic compound biosynthesis in plants and about their coupling in the secondary walls.

Different tentative conclusions can be considered from the current investigations in *bm1* maize. Several very different events are responsible for the *ZmCAD2-bm1* mutation in maize. The *ZmCAD2* gene encodes the predominant CAD for monolignol production, even though other CAD gene(s) could be simultaneously involved, or may have substitutive

activity. Comparing the biochemical traits in transposon tagged and disrupted mutant highlighted the relevance of the marker characteristics that were shown in CAD2 deficient plants, including aldehyde and atypical compounds and linkages. Even if further experiments need to be done, CAD2 is likely a good target for the improvement of energy production based on maize and grass lignocellulose biomass, including also a greater susceptibility to (environmentally friendly) pretreatments (Maehara et al, 2011; Wu et al, 2011). These specific properties are related to the presence of unusual compounds incorporated into lignins, inducing a modified structure of the polymer spread over carbohydrates in smaller domains. Despite the interest of *bm1* maize hybrids for cattle feeding has not yet been established, contrarily to the proven efficiency of *bm1* plants for bioenergy production, the breeding of maize *bm1* hybrids is likely promising because the *ZmCAD2* mutations seem to induce limited unfavorable consequences on other agronomical traits, even if opposite results could be observed with a few genetic backgrounds (Lorenz et al, 2009).

For the improvement of cell wall traits for both feeding and industrial uses, genetic targets should be considered in the whole set of genes involved in secondary wall biosynthesis. Genes involved in lignin production are upstream regulated by MYB and NAC transcription factors (Sonbol et al, 2009; Fornalé et al, 2010; Zhong et al, 2011; Gray et al, 2012). Ferulate driven cross-linkages are both dependent on ferulic acid (more probably of feruloyl-CoA) and arabinoxylan biosynthesis (Barrière et al, 2009; Hatfield and Marita, 2010; Jung and Phillips, 2010; Piston et al, 2010; Jung et al, 2011). Lignified tissue patterning is regulated during plant growth by members of several gene families, including members of the zinc finger and HD-ZIP families (Barrière et al, 2009). Search for transposon-tagging mutants with improved cell wall degradability should indeed be a relevant strategy for the discovery of genes with still unknown important role in secondary wall assembly (Vermeris et al, 2007; Jung and Phillips, 2010). Mutants and favorable alleles of all these genes have then to be considered as targets during maize (and grass) breeding for enhanced energy value. However, while plant breeding for animal feeding and bioenergy production both requested the knowledge of genetic mechanisms involved in secondary wall assembly, the breeding of bioenergy plants has also to take into account the lignin network susceptibility to mild-alkali pretreatments.

### Acknowledgements

Authors thank Marty Sachs, Director of the maize stock center for the supply of *bm1* mutants, Natalia de Leon and Jim Coors (University of Wisconsin) for the supply of A619bm1, and all three for their interest in this work. Authors also thank Thomas Lübbenstedt,

Hongjun Liu, and Paul Scott for discussions and interaction during investigations on the *bm7* mutation. Christiane Minault, Dominique Denoue and Pascal Vernoux were in charge of plant cropping at INRA Lusignan. Laurent Cézard and Frédéric Legée (INRA Versailles), and Corinne Melin (INRA Lusignan) were in charge of lignin content and composition analyses, and cell wall degradability estimates. Investigations on the Ev2210 *bm1* were funded by the MaizeWall project of the French genomic network Génoplante.

### References

Ali F, Scott P, Bakht J, Chen Y, Lübbenstedt T, 2010. Identification of novel brown midrib genes in maize by tests of allelism. *Plant Breeding* 129: 724-726

Aufrère J, Michalet-Doreau B, 1983. In vivo digestibility and prediction of digestibility of some by-products, pp. 25-33. In: EEC seminar, Melle Gontrode. 26-29 September

Barnes RF, Muller LD, Colenbrander VF, 1971. In vitro dry matter disappearance of brown-midrib mutants. *J Anim Sci* 33: 881-884

Barrière Y, Argillier O, 1993. Brown-midrib genes of maize: a review. *Agronomie* 13: 865-876

Barrière Y, Ralph J, Méchin V, Guillaumie S, Grabber JH, Argillier O, Chabbert B, Lapierre C, 2004a. Genetic and molecular basis of grass cell wall biosynthesis and degradability. II. Lessons from brown-midrib mutants. *Comptes Rendus Biologie* 327: 847-860

Barrière Y, Emile JC, Trainneau R, Surault F, Briand M and Gallais A, 2004b. Genetic variation for organic matter and cell wall digestibility in silage maize. Lessons from a 34-year long experiment with sheep in digestibility crates. *Maydica* 49: 115-126

Barrière Y, Méchin V, Lafarguette F, Manicacci D, Guillot F, Wang H, Laurensse D, Pichon M, Bosio M, Tatout C, 2009. Toward the discovery of maize cell wall genes involved in silage maize quality and capacity to biofuel production. *Maydica* 54: 161-198

Barrière Y, Charcosset A, Denoue D, Madur D, Baujand C, Laborde J, 2010. Genetic variation for lignin content and cell wall digestibility in early maize lines derived from ancient landraces. *Maydica* 55: 65-74

Baucher M, Chabbert B, Pilate G, Van Doorsselaere J, Tollier MT, Petit-Conil M, Cornu D, Monties B, Van Montagu M, Inze D, Jouanin L, Boerjan W, 1996. Red xylem and higher lignin extractability by down-regulating a cinnamyl alcohol dehydrogenase in poplar. *Plant Physiol* 112: 1479-1490

Burnham CR, 1947. Maize Genetics Cooperation Newsletter 21, pp 36

Burnham CR, Brinks RA, 1932. Linkage relations of a second brown-midrib gene (*bm2*) in maize. *J Am Soc Agron* 24: 960-963

Chavignau H, Goué N, Courtial A, Jouanin L, Rey-

mond M, Méchin V, Barrière Y, 2012. QTL for floral stem lignin content and degradability in three recombinant inbred line (RIL) progenies of *Arabidopsis thaliana* and search for candidate genes involved in cell wall biosynthesis and degradability. OJGen 2: 7-30

Chen W, Van Opdorp N, Fitzl D, Tewari J, Friedemann P, Greene T, Thompson S, Kumpatla S, Zheng P, 2012a. Transposon insertion in a cinnamyl alcohol dehydrogenase gene is responsible for a brown midrib1 mutation in maize. Plant Mol Biol 80: 289-397

Chen Y, Liu H, Ali F, Scott MP, Ji O, Frei UK, Lübbertedt T, 2012b. Genetic and physical fine mapping of the novel brown midrib gene *bm6* in maize (*Zea mays* L) to a 180 kb region on chromosome 2. Theor Appl Gen 25: 1223-1235

Corpet F, 1988. Multiple sequence alignment with hierarchical clustering. Nucl Acids Res 16: 10881-10890

Corredor DY, Salazar JM, Hohn KL, Bean S, Bean B, Wang D, 2009. Evaluation and characterization of forage sorghum as feedstock for fermentable sugar production. Appl Biochem Biotechnol 158: 164-179

Damiani I, Morreel K, Danoun S, Goeminne G, Yahiaoui N, Marque C, Kopka J, Messens E, Goffner D, Boerjan W, Boudet AM, Rochange S, 2005. Metabolite profiling reveals a role for atypical cinnamyl alcohol dehydrogenase CAD1 in the synthesis of coniferyl alcohol in tobacco xylem. Plant Mol Biol 59: 753-769

Dence CW, Lin SY, 1992. The determination of lignin, pp. 33-61. In: Methods in lignin chemistry. Lin SY, Dence CW eds. Springer-Verlag, Berlin, Germany

Dien BS, Sarah G, Pedersen JF, Sattler SE, Chen H, Funnell-Harris DL, Nichols NN, Cotta MA, 2009. Improved Sugar Conversion and ethanol yield for forage sorghum (*Sorghum bicolor* L. Moench) lines with reduced lignin contents. Bionergy Res 2: 153-164

Dolstra O, Medema JH, 1990. An effective screening method for genetic improvement of cell-wall digestibility in forage maize, pp. 258-270. In: Proceedings 15th congress maize and sorghum section of Eucarpia. 4-8 June 1990, Baden, Austria

Emerson RA, 1935. Cornell Univ Agric Exp Stn, Memoir 180

Eyster WH. 1926 Chromosome VIII in maize Science 64: 22

Fornalé S, Shi X, Chai C, Encina A, Irar S, Capellades M, Fuguet E, Torres JL, Rovira P, Puigdomènec P, Rigau J, Grotewold E, Gray J, Caparrós-Ruiz D, 2010. ZmMYB31 directly represses maize lignin genes and redirects the phenylpropanoid metabolic flux. Plant J 64: 633-644

Fornalé S, Capellades M, Encina A, Wang K, Irar S, Lapierre C, Ruel K, Joseleau JP, Berenguer J, Puigdomènec P, Rigau J, Caparrós-Ruiz D, 2012. Altered lignin biosynthesis improves cellulosic bioethanol production in transgenic maize plants down-regulated for cinnamyl alcohol dehydrogenase. Mol Plant 5: 817-830

Gee MS, Nelson OE, Kuc J, 1968. The abnormal lignins produced by the brown-midrib mutants of maize. II. Comparative studies on normal and brown-midrib-1 dimethylfornamide lignins. Arch Biochem Biophys 123: 403-408

Gill GP, Brown GR, Neale DB, 2003. A sequence mutation in the cinnamyl alcohol dehydrogenase gene associated with altered lignification in loblolly pine. Plant Biotechnol J 1: 253-258

Goering HK, van Soest PJ, 1970. Forage fiber analysis (Apparatus, reagents, procedures and some applications), pp. 1-379. US Dept Agri Sc. Handbook

Goffner D, Joffroy I, Grima-Pettenati J, Halpin C, Knight ME, Schuch W, Boudet AM, 1992. Purification and characterization characterization of isoforms of cinnamyl alcohol dehydrogenase (CAD) from Eucalyptus xylem. Planta 188: 48-53

Goffner D, van Doorselaere J, Yahiaoui N, Samaj J, Grima-Pettenati J, Boudet AM, 1998. A novel aromatic alcohol dehydrogenase in higher plants: molecular cloning and expression. Plant Mol Biol 36: 755-765

Gray J, Caparros-Ruiz D, Grotewold E, 2012. Grass phenylpropanoids: Regulate before using! Plant Science 184: 112-120

Guillaumie S, Pichon M, Martinant JP, Bosio M, Goffner D, Barrière Y, 2007a. Differential expression of phenylpropanoid and related genes in brown-midrib *bm1*, *bm2*, *bm3*, and *bm4* young near-isogenic maize plants. Planta 226: 235-250

Guillaumie S, San-Clemente H, Deswarte C, Martinez Y, Lapierre C, Murigneux A, Barrière Y, Pichon M, Goffner D, 2007b. MAIZEWALL. Database and developmental gene expression profiling of cell wall biosynthesis and assembly in maize. Plant Physiol 143: 339-363

Halpin C, Holt K, Chojecki J, Oliver D, Chabbert B, Monties B, Edwards K, Barakate A, Foxon GA, 1998. Brown-midrib maize (*bm1*) - a mutation affecting the cinnamyl alcohol dehydrogenase gene. Plant J 14: 545-553

Haney LJ, Hake S, Scott MP, 2008. Allelism testing of Maize Coop Stock Center lines containing unknown brown midrib alleles. Maize Genetics Cooperative Newsletter 82: 4-5

Hatfield RD, Jung HJG, Ralph J, Buxton DR, Weimer PJ, 1994. A comparison of the insoluble residues produced by the Klason lignin and acid detergent lignin procedures. J Sci Food Agric 65: 51-58

Hatfield RD, Fukushima RS, 2005. Can lignin be accurately measured. Crop Sci. 45: 832-839

Hatfield RD, Marita JM, 2010. Enzymatic processes involved in the incorporation of hydroxycinnamates into grass cell walls. Phytochem Rev 9:

35-45

Hawkins SW, Boudet AM, 1994. Purification and characterization of cinnamyl alcohol-dehydrogenase isoforms from the periderm of *Eucalyptus gunnii* Hook. *Plant Physiol* 104: 75-84

Jacquet G, 1997. Structure et réactivité des lignines de graminées et des acides phénoliques associés: développement des méthodologies d'investigation. Ph.D. thesis, Université Aix-Marseille III

Jorgenson LR, 1931. Brown midrib in maize and its lineage relations. *J Am Soc Agron* 23:549-557

Jornvall H, Persson B, Jeffery J, 1987, Characteristics of alcohol/polyol dehydrogenases. The zinc-containing long-chain alcohol dehydrogenases. *Eur J Biochem* 167: 195-201

Jornvall H, Persson B, Krook M, Atrian S, Gonzalez-duarte R, Jeffery J, Ghosh D, 1995. Short-chain dehydrogenases reductases (Sdr). *Biochemistry* 34: 6003-6013

Jung H, Mertens D, Payne A, 1997. Correlation of acid detergent lignin and Klason lignin with digestibility of forage dry matter and neutral detergent fiber. *J Dairy Sci* 80: 1622-1628

Jung HG, Phillips RL, 2010. Putative seedling ferulate ester (sfe) maize mutant: morphology, biomass yield, and stover cell wall composition and rumen degradability. *Crop Sci* 50: 403-418

Jung HG, Mertens D, Phillips RL, 2011. Effect of reduced ferulate-mediated lignin/arabinoxylan cross-linking in corn silage on feed intake, digestibility, and milk production. *J Dairy Sci* 94: 5124-5137

Kanazawa K, Goodman MM, O'Malley DM, 1999. Genetic and biochemical analysis of maize CAD. January 17-21. Town & Country Hotel, San Diego, USA

Kiesselsbach TA, 1922. Corn investigations. *Nebr Agric Exp Sta Res Bull* 20: 51-52

Kim H, Ralph J, Lu F, Pilate G, Leplé JC, Pollet B, Lapierre C, 2002. Identification of the structure and origin of thioacidolysis marker compounds for cinnamyl alcohol dehydrogenase deficiency in angiosperms. *J Biol Chem* 277: 47412-47419

Kim H, Ralph J, Lu F, Ralph SA, Boudet AM, MacKay JJ, Sederoff RR, Ito T, Kawai S, Ohashi H, Higuchi T, 2003. NMR analysis of lignins in CAD-deficient plants. Part 1. Incorporation of hydroxycinnamaldehydes and hydroxybenzaldehydes into lignins. *Organic Biomolecular Chemistry* 1: 158-281

Kuc J, Nelson OE, 1964. The abnormal lignins produced by the brown-midrib mutants of maize. I. The *brown-midrib-1* mutant. *Arch Biochem Biophys* 105: 103-113

Lapierre C, Monties B, Rolando C, 1988. Thioacidolyses of diazomethane-methylated pine compression wood and wheat straw *in situ* lignins. *Holzforschung* 42: 409-411

Lapierre C, 1993. Applications of new methods for the investigation of lignin structure, pp. 133-136. In: *Forage Cell Wall Structure and Digestibility*. Jung HG, Buxton DR, Hatfield RD eds. American ASA-CSSA-SSSA, Madison, Wisconsin, USA

Lapierre C, Pollet B, Petit-Conil M, Toval G, Romero J, Pilate G, Leple JC, Boerjan W, Ferret V V, De Nadai V, Jouanin L, 1999. Structural alterations of lignins in transgenic poplars with depressed cinnamyl alcohol dehydrogenase or caffeic acid O-methyltransferase activity have an opposite impact on the efficiency of industrial kraft pulping. *Plant Physiol* 119: 153-164

Lapierre C, Pollet B, MacKay JJ, Sederoff RR, 2000. Lignin structure in a mutant pine deficient in cinnamyl alcohol dehydrogenase. *J Agric Food Chem* 48: 2326-2331

Lapierre C, Pilate G, Pollet B, Mila I, Leplé JC, Jouanin L, Kim H, Ralph J, 2004. Signatures of cinnamyl alcohol dehydrogenase deficiency in poplar lignins. *Phytochem* 65: 313-321

Lapierre C, 2010. Determining lignin structure by chemical degradations. In *Lignin and lignans - Advances in Chemistry*, C. Heitner, D. Dimmel, and J.A. Schmidt, eds (Boca Raton, USA: CRC Press, Taylor & Francis Group), pp 11-48

Lechtenberg VL, Muller LD, Bauman LF, Rhykerd CL, Barnes RF, 1972. Laboratory and *in vivo* evaluation of inbred and  $F_2$  populations of brown-midrib mutants of *Zea mays* L. *Agron J* 64: 657-680

Lorenz AJ, Coors JG, de Leon N, Wolfrum EJ, Hames BR, Sluiter AD, Weimer PJ, 2007. Characterization, genetic variation, and combining ability of maize traits relevant to the production of cellulosic ethanol. *Crop Sci* 49:85-98

Maehara T, Takai T, Ishihara H, Yoshida M, Fukuda K, Gau M, Kaneko S, 2011. Effect of lime pretreatment of brown midrib sorghum. *Biosci Biotechnol Biochem* 75: 2415-2417

Mir Derikvand M, Berrio Sierra J, Ruel K, Pollet B, Do CT, Thévenin J, Buffard D, Jouanin L, Lapierre C, 2008. Redirection of the phenylpropanoid pathway to feruloyl malate in *Arabidopsis* mutants deficient for cinnamoyl-CoA reductase 1. *Planta* 227: 943-956

Morrow SL, Mascia P, Self KA, Altschuler M, 1997. Molecular characterization of a brown midrib3 deletion mutation in maize. *Molecular Breeding* 3: 351-357

O'Connell A, Holt K, Piquemal J, Grima-Pettenati J, Boudet A, Pollet B, Lapierre C, Petit-Conil M, Schuch W, Halpin C, 2002. Improved paper pulp from plants with suppressed cinnamoyl-CoA reductase or cinnamyl alcohol dehydrogenase. *Transgenic Res* 11: 495-503

Oliver AL, Grant RJ, Pedersen JF, O'Rear J, 2003. Comparison of brown midrib-6 and -18 forage sorghum with conventional sorghum and corn silage in diets of lactating dairy cows. *J Dairy Sci* 87: 637-644

Pillonel C, Mulder MM, Boon JJ, Forster B, Binder A, 1991. Involvement of cinnamyl-alcohol dehydrogenase in the control of lignin formation in *Sorghum bicolor* L. Moench. *Planta* 185: 538-544

Pandey B, Pandey VP, Dwivedi UN, 2011. Cloning, expression, functional validation and modeling of cinnamyl alcohol dehydrogenase isolated from xylem of *Leucaena leucocephala*. *Protein Expr Purif* 79: 197-203

Provan GJ, Scobbie L, Chesson A, 1997. Characterisation of lignin from CAD and OMT deficient *bm* mutant of maize. *J Agric Food* 73: 133-142

Ralph J, Hatfield RD, Quideau S, Helm RF, Grabber JH, Jung HJG, 1994. Pathway of *p*-coumaric acid incorporation into maize lignin as revealed by NMR. *J Am Chem Soc* 116: 9448-9456

Ralph J, Kim H, Lu F, Grabber J, Leplé JC, Berrio-Sierra J, Mir Derikvand M, Jouanin L, Boerjan W, Lapierre C, 2008. Identification of the structure and origin of a thioacidolysis marker compound for ferulic acid incorporation into angiosperm lignins (and an indicator for Cinnamoyl-CoA Reductase deficiency). *Plant J* 53: 368-379

Riboulet C, Guillaumie S, Méchin V, Bosio M, Pichon M, Goffner D, Lapierre C, Pollet B, Lefèvre B, Martinant JP, Barrière Y, 2009. Kinetics of phenylpropanoid gene expression in maize growing internodes: Relationships with cell wall deposition. *Crop Sci* 49: 211-223

Saathoff AJ, Sarath G, Chow EK, Dien BS, Tobias CM, 2011. Down-regulation of cinnamyl-alcohol dehydrogenase in switchgrass by RNA silencing results in enhanced glucose release after cellulase treatment. *PLoS One* 2011 6(1): e16416

Saathoff AJ, Hargrove MS, Haas EJ, Tobias CM, Twigg P, Sattler S, Sarath G, 2012. Switchgrass PviCAD1: Understanding residues important for substrate preferences and activity. *Appl Biochem Biotechnol* 168: 1086-1100

Saballos A, Vermerris W, Rivera L, Ejeta G, 2008. Allelic association, chemical characterization and saccharification properties of brown midrib mutants of sorghum (*Sorghum bicolor* (L) Moench). *Bioenerg Res* 1: 193-204

Saballos A, Ejeta G, Sanchez E, Kang C, Vermerris W, 2009. A genomewide analysis of the cinnamyl alcohol dehydrogenase family in sorghum [*Sorghum bicolor* (L) Moench] identifies *SbCAD2* as the brown midrib6 gene. *Genetics* 181: 783-795

Sattler SE, Saathoff AJ, Haas EJ, Palmer NA, Funnell-Harris DL, Sarath G, Pedersen JF, 2009. A non-sense mutation in a cinnamyl alcohol dehydrogenase gene is responsible for the Sorghum brown midrib6 phenotype. *Plant Physiol.* 150: 584-595.

Sattler SE, Funnell-Harris DL, Pedersen JF, 2010a. Efficacy of singular and stacked brown midrib 6 and 12 in the modification of lignocellulose and grain chemistry. *J Agric Food Chem* 58: 3611-3616

Sattler SE, DL Funnell-Harris, JF Pedersen, 2010b. Brown midrib mutations and their importance to the utilization of maize, sorghum, and pearl millet lignocellulosic tissues. *Plant Science* 178: 229-238

Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, Liang C, Zhang J, Fulton L, Graves TA, Minx P, Reily AD, Courtney L, Kruchowski SS, Tomlinson C, Strong C, Delehaunty K, Fronick C, Courtney B, Rock SM, Belter E, Du F, Kim K, Abbott RM, Cotton M, Levy A, Marchetto P, Ochoa K, Jackson SM, Gillam B, Chen W, Yan L, Higginbotham J, Cardenas M, Waligorski J, Applebaum E, Phelps L, Falcone J, Kanchi K, Thane T, Scimone A, Thane N, Henke J, Wang T, Ruppert J, Shah N, Rotter K, Hodges J, Ingenthron E, Cordes M, Kohlberg S, Sgro J, Delgado B, Mead K, Chinwalla A, Leonard S, Crouse K, Collura K, Kudrna D, Currie J, He R, Angelova A, Rajasekar S, Mueller T, Lomeli R, Scara G, Ko A, Delaney K, Wissotski M, Lopez G, Campos D, Braidotti M, Ashley E, Golser W, Kim H, Lee S, Lin J, Dujmic Z, Kim W, Talag J, Zuccolo A, Fan C, Sebastian A, Kramer M, Spiegel L, Nascimento L, Zutavern T, Miller B, Ambroise C, Muller S, Spooner W, Narachania A, Ren L, Wei S, Kumari S, Faga B, Levy MJ, McMahan L, Van Buren P, Vaughn M, Ying K, Yeh C-T, Emrich S, Jia Y, Kalyanaraman A, Hsia A-P, Barbazuk WB, Baucom RS, Brutnell TP, Carpita NC, Chaparro C, Chia J-M, Deragon J-M, Estill JC, Fu Y, Jeddelloh JA, Han Y, Lee H, Li P, Lisch DR, Liu S, Liu Z, Nagel DH, McCann MC, SanMiguel P, Myers AM, Nettleton D, Nguyen J, Penning BW, Ponnala L, Schneider KL, Schwartz DC, Sharma A, Soderlund C, Springer NM, Sun Q, Wang H, Waterman M, Westerman R, Wolfgruber TK, Yang L, Yu Y, Zhang L, Zhou S, Zhu Q, Bennetzen JL, Dawe RK, Jiang J, Jiang N, Presting GG, Wessler SR, Aluru S, Martienssen RA, Clifton SW, McCombie WR, Wing RA, Wilson RK, 2009. The B73 maize genome: complexity, diversity, and dynamics. *Science* 326: 1112-1115

Scott PM, Sattler SE, Funnell-Harris DE, Pedersen JF, 2010. Brown midrib mutations and their importance to the utilization of maize, sorghum, and pearl millet lignocellulosic tissues. *Plant Science* 178: 229-238

Shi C, Koch G, Ouzunova M, Wenzel G, Zein I, Lübbertedt T, 2006. Comparison of maize brown-midrib isogenic lines by cellular UV-microspectrophotometry and comparative transcript profiling. *Plant Mol Biol* 62: 697-714

Sibout R, Eudes A, Mouille G, Pollet B, Lapierre C, Jouanin L, Séguin A, 2005. CINNAMYL ALCOHOL DEHYDROGENASE-C and -D are the primary genes involved in lignin biosynthesis in the floral stem of *Arabidopsis*. *Plant Cell* 17: 2059-2076

Sonbol FM, Fornalé S, Cappellades M, Encina A, Tourino S, Torres JL, Rovira P, Ruel K, Puig-

domenech P, Rigau J, Caparros-Ruiz D, 2009. The maize ZmMYB42 represses the phenylpropanoid pathway and affects the cell wall structure, composition and degradability in *Arabidopsis thaliana*. *Plant Mol Biol* 70: 283-296

Struik P. 1983. Physiology of forage maize (*Zea mays* L) in relation to its production and quality. PhD Dissertation, Agricultural University, 6700 GW Wageningen, The Netherlands

Thévenin J, Pollet B, Letarne B, Saulnier L, Gissot L, Maia-Grondard A, Lapierre C, Jouanin L. 2011. The simultaneous repression of CCR and CAD, two enzymes of the lignin biosynthetic pathway, results in sterility and dwarfism in *Arabidopsis thaliana*. *Mol Plant* 4: 70-82

Vanholme R, Morreel K, Darrah C, Oyarce P, Grabber JH, Ralph J, Boerjan W, 2012. Metabolic engineering of novel lignin in biomass crops. *New Phytol* 196: 978-1000

Vermerris W, Saballos A, Ejeta G, Mosier NS, Ladish MR, Carpita NC, 2007. Molecular breeding to enhance ethanol production from corn and sorghum stover. *Crop Sci* 47: S142-S153

Vignols F, Rigau J, Torres MA, Capellades M, Puigdomenech P, 1995. The *brown midrib 3* (*bm3*) mutation in maize occurs in the gene encoding caffeic acid O-methyltransferase. *Plant Cell* 7: 407-416

Wu L, Mitsuhiro A, Masakazu I, Masahisa W, Takai T, Gau M, Tokuyasu K, 2011. Low temperature alkali pretreatment for improving enzymatic digestibility of sweet sorghum bagasse for ethanol production. *Bioresource Technol* 102: 4793-4799

Zhang K, Qian Q, Huang Z, Wang Y, Li M, Hong L, Zeng D, Gu M, Chu C, Cheng Z. 2006. GOLD HULL AND INTERNODE2 encodes a primarily multifunctional cinnamyl-alcohol dehydrogenase in rice. *Plant Physiol* 140: 972-983

Zhang Y, Culhaoglu T, Pollet B, Melin C, Denoue D, Barrière Y, Baumberger S, Méchin V, 2011. Impact of lignin structure and cell wall reticulation on maize cell wall degradability. *J Sci Food Agric* 59:10129-10135

Zhong R, Lee C, McCarthy RL, Reeves CK, Jones EG, Ye ZH, 2011. Transcriptional activation of secondary wall biosynthesis by rice and maize NAC and MYB transcription factors. *Plant Cell Physiol* 52: 1856-1871

**Supplementary Figure 1** - Nucleotide alignments of the ZmCAD2 CDS in normal and bm1 isogenic lines (Identical nucleotides are shaded in grey, X are missing data)

Ev2210	GCTGGACTAC	ATCATCGACA	CGGTGCCCGT	GCACCACCCG	CTGGAGCCGT	ACCTGGCGCT	GCTGAAGCTG	GACGGCAAGC	TCGTGCTGCT	GGCGCTCATC
F2bml	GCTGGACTAC	ATCATCGACA	CGGTGCCCGT	GCACCACCCG	CTGGAGCCGT	ACCTGGCGCT	GCTGAAGCTG	GACGGCAAGC	TCGTGCTGCT	GGCGCTCATC
511Jbml	GCTGGACTAC	ATCATCGACA	CGGTGCCCGT	GCACCACCCG	CTGGAGCCGT	ACCTGGCGCT	GCTGAAGCTG	GACGGCAAGC	TCGTGCTGCT	GGCGCTCATC
511Kbml	GCTGGACTAC	ATCATCGACA	CGGTGCCCGT	GCACCACCCG	CTGGAGCCGT	ACCTGGCGCT	GCTGAAGCTG	GACGGCAAGC	TCGTGCTGCT	GGCGCTCATC
5803Cbml	GCTGGACTAC	ATCATCGACA	CGGTGCCCGT	GCACCACCCG	CTGGAGCCGT	ACCTGGCGCT	GCTGAAGCTG	GACGGCAAGC	TCGTGCTGCT	GGCGCTCATC
5803Ibml7	GCTGGACTAC	ATCATCGACA	CGGTGCCCGT	GCACCACCCG	CTGGAGCCGT	ACCTGGCGCT	GCTGAAGCTG	GACGGCAAGC	TCGTGCTGCT	GGCGCTCATC
F7803Cbml	GCTGGACTAC	ATCATCGACA	CGGTGCCCGT	GCACCACCCG	CTGGAGCCGT	ACCTGGCGCT	GCTGAAGCTG	GACGGCAAGC	TCGTGCTGCT	GGCGCTCATC
Ev2210bm1	GCTGGACTAC	ATCATCGACA	CGGTGCCCGT	GCACCACCCG	CTGGAGCCGT	ACCTGGCGCT	GCTGAAGCTG	GACGGCAAGC	TCGTGCTGCT	GGCGCTCATC

	901	1000								
B73	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
F2	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTAG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
Ev2210	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
F2bml	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
511Jbml	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
511Kbml	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
5803Cbml	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
5803Ibml7	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
F7803Cbml	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC
Ev2210bm1	GGCGAGCCCC	TGAGCTTCGT	GTGCGCCATG	GTGATGCTGG	GGCGGAAGGC	CATCACGGGG	AGCTTCATCG	GCAGCATCGA	CGAGACCGCT	GAGGTGCTTC

	1001	1100								
B73	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
F2	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
Ev2210	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
F2bml	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
511Jbml	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
511Kbml	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
5803Cbml	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
5803Ibml7	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
F7803Cbml	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA
Ev2210bm1	AGTTCTGGT	CGACAAGGGG	CTCACCTCCC	AGATCGAGGT	GGTCAAGATG	GGGTACGTGA	ACGAGGCGCT	GGAGCGGCTG	GAGCGCAACG	ACGTCCGCTA

	1101	1173							
B73	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
F2	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
Ev2210	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
F2bml	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
511Jbml	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
511Kbml	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
5803Cbml	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
5803Ibml7	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
F7803Cbml	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	
Ev2210bm1	CCGCTTCGTC	GTCGACGCTG	CCGGTAGCAA	CGTCGAGGCC	GAGGCGGCCG	GGCGGGATGC	GGCCAGCAAC	TGA	

**Supplementary Figure 2 - Alignments of the deduced amino-acid sequences of the ZmCAD2 protein in all investigated lines (Identical amino acid residues are shaded in grey, X are missing data).**

	1	80
B73	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
F2	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
Ev2210	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
F2bml	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
511Jbml	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
511Kbml	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
5803Cbm1	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
5803Ibm7	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
F7803bml	MGSLASERKV VGWAARDATG HLSPYSYTLX XXXXXXXXXXXX XXXXXXXXXXXX XXXXXXXXXXXX XXXXXXXXXXXHE VVGEVVVEVGP	
Ev2210bml	MGSLASERKV VGWAARDATG HLSPYSYTLR NTGPEDVVVK VLYCGICHTD IHQAKNHLGA SKYPMVPGHE VVGEVVVEVGP	
	81	160
B73	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGGFASTMV VDQKFVVKIP AGLAPEQAAAP	
F2	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGGFASTMV VDQKFVVKIP AGLAPEQAAAP	
Ev2210	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGGFASTMV VDQKFVVKIP AGLAPEQAAAP	
F2bml	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGGFASTMV VDQKFVVKIP AGLAPEQAAAP	
511Jbml	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNARLH .....	
511Kbml	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGEFASTMV VDQKFVVKIP AGLAPEQAAAP	
5803Cbm1	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGGFACTMV VDQKFVVKIP AGLAPEQAAAP	
5803Ibm7	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGGFACTMV VDQKFVVKIP AGLAPEQAAAP	
F7803bml	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGGFASTMV VDQKFVVKIP AGLAPEQAAAP	
Ev2210bml	EVAKYGVGDV VGVGVIVGCC RECPCKANV EQYCNKKIWS YNDVYTDGRP TQGGFASTMV VDQKFVVKIP AGLAPEQAAAP	
	161	240
B73	LLCAGTVTVS PLKHFGLLTP GLRGGILGLG GVGHMGVKVA KAMGHHVTVI SSSSKKRAEA MDHLGADAYL VSSDAAAMAA	
F2	LLCAGTVTVS PLKHFGLLTP GLRGGILGLG GVGHMGVKVA KAMGHHVTVI SSSSKKRAEA MDHLGADAYL VSSDAAAMAA	
Ev2210	LLCAGTVTVS PLKHFGLLTP GLRGGILGLG GVGHMGVKVA KAMGHHVTVI SSSSKKRAEA MDHLGADAYL VSSDAAAMAA	
F2bml	.....	
511Jbml	.....	
511Kbml	LLCAGTVTVS PLKHFGLLTP GLRGGILGLG GVGHMGVKVA KAMGHHVTVI SSSSKKRAEA MDHLGADAYL VSSDAAAMAA	
5803Cbm1	LLCAGTVTVS PLKHFGLLTP GLRGGILGLG GVGHMGVKVA KAMGHHVTVI SSSSKKRAEA MDHLGADAYL VSSDAAAMAA	
5803Ibm7	LLCAGTVTVS PLKHFGLLTP GLRGGILGLG GVGHMGVKVA KAMGHHVTVI SSSSKKRAEA MDHLGADAYL VSSDAAAMAA	
F7803bml	LLCAGTVTVS PLKHFGLLTP GLRGGILGLG GVGHMGVKVA KAMGHHVTVI SSSSKKRAEA MDHLGADAYL VSSDAAAMAA	
Ev2210bml	LLCAGTVTVS PLKHFGLLTP GLRGGILGLG GVGHMGVKVA KAMGHHVTVI SSSSKKRAEA MDHLGAR .....	
	241	320
B73	AADSLDYIID TVPVHHPLEP YLALLKLDGK LVLLGVIGEP LSFVSPMVML GRKAITGSFI GSIDETAEVL QFCVDKGLTS	
F2	AADSLDYIID TVPVHHPLEP YLALLKLDGK LVLLGVIGEP LSFVSPMVML GRKAITGSFI GSIDETAEVL QFCVDKGLTS	
Ev2210	AADSLDYIID TVPVHHPLEP YLALLKLDGK LVLLGVIGEP LSFVSPMVML GRKAITGSFI GSIDETAEVL QFCVDKGLTS	
F2bml	.....	
511Jbml	.....	
511Kbml	AADSLDYIID TVPVHHPLEP YLALLKLDGK LVLLGVIGEP LSFVSPMVML GRKAITGSFI GSIDETAEVL QFCVDKGLTS	
5803Cbm1	AADSLDYIID TVPVHHPLEP YLALLKLDGK LVLLGVIGEP LSFVSPMVML GRKAITGSFI GSIDETAEVL QFCVDKGLTS	
5803Ibm7	AADSLDYIID TVPVHHPLEP YLALLKLDGK LVLLGVIGEP LSFVSPMVML GRKAITGSFI GSIDETAEVL QFCVDKGLTS	
F7803bml	AADSLDYIID TVPVHHPLEP YLALLKLDGK LVLLGVIGEP LSFVSPMVML GRKAITGSFI GSIDETADVL QFCVDKGLTS	
Ev2210bml	.....	
	321	367
B73	QIEVVKMGYV NEALERLERN DVRYRFFVVDV AGSNVEAEAA AAAAASN	
F2	QIEVVKMGYV NEALERLERN DVRYRFFVNV AGSNVEAEAA AAAAASN	
Ev2210	QIEVVKMGYV NEALERLERN DVRYRFFVVDV AGSNVEAEAA AAAAASN	
F2bml	.....	
511Jbml	.....	
511Kbml	QIEVVKMGYV NEALERLERN DVRYRFFVVDV AGSNVEAEAA AAAAASN	
5803Cbm1	QIEVVKMGYV NEALERLERN DVRYRFFVVDV AGSNVEAEAA AAAAASN	
5803Ibm7	QIEVVKMGYV NEALERLERN DVRYRFFVVDV AGSNVEAEAA AAAAASN	
F7803bml	QIEVVKMGYV NEALERLERN DVRYRFFVVDV AGSNVEAEAA AAAAASN	
Ev2210bml	.....	