

CRISTINA CASTRACANI (*) - FIORENZA A. SPOTTI (*) - DONATO A. GRASSO (*)
ALBERTO FANFANI (**) - ALESSANDRA MORI (*)

A NEW EXCEPTION TO THE DOMINANCE-DISCOVERY TRADE-OFF RULE IN ANT COMMUNITIES⁽¹⁾

(*) Dipartimento di Bioscienze, Università degli Studi di Parma, Parco delle Scienze 11/A, 43124 Parma, Italy; e-mail: cristina.castracani@nemo.it

(**) Dipartimento di Biologia e Biotecnologie 'Charles Darwin', Università degli Studi di Roma 'La Sapienza', Viale dell'Università 32, 00185 Roma, Italy.

Castracani C., Spotti F.A., Grasso D.A., Fanfani A., Mori A. - A new exception to the dominance-discovery trade-off rule in ant communities.

Interspecific competition is one of the main factor shaping ant community structure. The "dominance-discovery trade-off", an inverse relationship between interference and exploitative competitive ability, is thought to be a fundamental mechanism facilitating coexistence of ants. Despite its widely acceptance in literature, this trade-off appears to be broken in several cases and needs further investigations. Thus, the aim of the present study was to detect the presence/absence of the dominance-discovery trade-off in Mediterranean ant communities. Samplings were conducted inside the Presidential Estate of Castelporziano (Rome, Italy) and seven sampling sites were selected according to their vegetation structure. In each site, baiting was conducted to assess two behavioural indices for each species: the monopolization index and the discovery ability index. Correlation analyses showed no evidence for the existence of a dominance-discovery trade-off. Furthermore, a positive correlation between the two indices was detected in some of the sampling sites. The dominance-discovery trade-off seems therefore to be just one of the possible mechanisms facilitating species coexistence in ant communities.

KEY WORDS: interspecific competition, behavioural dominance, coexistence, ants, Castelporziano Estate.

INTRODUCTION

HÖLLOBLER and WILSON (1990) defined competition as "the hallmark of ant ecology". Actually, several studies pointed out that interspecific competition is one of the main factor shaping ant community structure (FELLERS, 1987; SAVOLAINEN and VEPSÄLÄINEN, 1988; HERBERS, 1989; ANDERSEN, 1992; SANDERS and GORDON, 2003; PARR *et al.*, 2005). As a consequence, a key issue becomes what allows multiple ant species to coexist in a given habitat (ANDERSEN, 2008; LESSARD *et al.*, 2009). One consolidated hypothesis suggests that an evolutionary trade-off between interference and exploitative competitive ability may facilitate coexistence of ants: this is called "the dominance-discovery trade-off" (FELLERS, 1987). In other words, the ability of a species to discover food resources is inversely related to its ability to defend them. Hence, behaviourally subordinate species, better at discovering food resources, may coexist with behaviourally dominant species that easily monopolise those resources. This trade-off is widely cited in the ant ecology literature and has consequently received general acceptance (SAVOLAINEN and VEPSÄLÄINEN, 1988; PERFECTO, 1994; FEENER, 2000; ADLER *et al.*, 2007; DELSINNE *et al.*, 2007; PARR and GIBB, 2010, 2012; PEARCE-DUVET and FEENER, 2010). On the other hand, several studies document how

the dominance-discovery trade-off can be broken by particular biotic factors (DAVIDSON, 1998; HOLWAY, 1999; LEBRUN and FEENER, 2007; ARNAN *et al.*, 2011) or environmental conditions (CERDÀ *et al.*, 1997; BESTELMEYER, 2000; SANTINI *et al.*, 2007; GIBB and PARR, 2010). Thus, the assumption that the trade-off is widespread in ant communities appears actually a hypothesis to be demonstrated and demands further investigations (PARR and GIBB, 2012). According to these requirements, the aim of the present study was to detect the presence/absence of the discovery-dominance trade-off in Mediterranean ant communities.

MATERIALS AND METHODS

Samplings were conducted inside the Presidential Estate of Castelporziano (Rome, Italy), which covers an area of almost 6000 hectares between Rome (S-SW) and the Tyrrhenian cost. Originally used as hunting reserve and farm, this area progressively changed its functions: since 1977 the hunting activity has been prohibited and the cultivated fields offer now a good example of eco-compatible management of agro-ecosystems. Declared Natural Reserve in 1999, the Estate shows a wide range of Mediterranean ecosystems and high levels of biodiversity: it hosts more than 1000 species of plants and more than 2900 species of animals. The area of the Natural Reserve is between 0 and 85 m (a.m.s.l.), with a mean slope of 5% to the sea, and the climate is meso-Mediterranean with a hot and dry period in summer (AA.VV., 2001, 2006).

Seven sampling sites were selected to study the most representative habitats of the Estate (DELLA ROCCA *et al.*,

¹ Original scientific contribution presented and discussed at XV Meeting of A.I.S.A.S.P., Italian Section of I.U.S.S.I. (International Union for the Study of Social Insects); Reggio Emilia - Italy, 18-19 September 2014.

- 2001). They are described as follows (for further details, see CASTRACANI *et al.*, 2010; SPOTTI *et al.*, 2010):
1. Cultivated Field (CF) - Fields cultivated with lucerne (*Medicago sativa*)
 2. Dry Grassland (DG) - Dry pasture lands (*Dasyphyrum villosum* and *Vulpia ligustica*)
 3. Wet Grassland (WG) - Wet pasture lands (Molinio-Arrhenatheretea group plants)
 4. Pine-wood (PW) - Reforestations with umbrella pines (*Pinus pinea*)
 5. Pure Ilex Grove (PIG) - High scrub and Mediterranean wood (*Quercus ilex*)
 6. Mixed Oak Wood (MOW) - Tyrrhenian deciduous oak wood (*Quercus cerris*, *Quercus frainetto*, *Carpinus orientalis*)
 7. Mediterranean Scrub (MS) - Acidophilous Mediterranean woods and bushes (*Erica arborea*, *Quercus suber* and, subordinately, *Quercus ilex*).

The first three sites (CF, DG, WG) are characterized by the presence of low vegetation, whereas the other four sites (PW, PIG, MOW, MS) are arboreal habitats (high vegetation sites).

Sampling took place during July 2007. In every site, three parallel 100-m linear transects were set up: each one consisting of 10 traps with 10-m spacing. The first transect was composed of pitfall traps: the experimental protocol and the results from this data collection are shown in SPOTTI *et al.* (2010). The other two transects consisted of two types of food baits: cookie + honey (10 baits) or tuna + oil (10 baits). All resources were placed on graph paper squares (7x7 cm) with scotch-tape on the lower side to prevent liquid leakage under the bait. This kind of baits are commonly used in studies on ant community ecology to simulate naturally occurring, patchy, large food resources for which ants frequently compete (FELLERS, 1987; ANDERSEN, 1992; CERDÀ *et al.*, 1997; AGOSTI *et al.*, 2000; PARR *et al.*, 2005; FEENER *et al.*, 2008; LESSARD *et al.*, 2009; STUBLE *et al.*, 2012). Each bait was observed at 5, 15, 30 and 60 minutes after the beginning of the experiment. For each species, the number of individuals feeding on the resource was recorded. Ants at baits were identified directly in the field or collected for further identification using the dichotomous keys of HÖLLOBLER and WILSON (1990) and BOLTON (1994). For some specimens identification at species level was achieved with the collaboration of Dr. Fabrizio Rigato (Natural History Museum of Milan, Italy). All specimens are preserved in the Laboratory of Myrmecology, Department of Life Sciences, University of Parma, Italy. In order to cover a wider range of foraging activity, baiting was run in early morning and replicated in late afternoon. A total of 280 baits were placed (10 baits x 2 types of food x 2 periods of the day x 7 site types).

For each species, data from baits enabled to calculate two behavioural indices: the Monopolisation Index (MI) and the Discovery Ability Index (DAI) (SANTINI *et al.*, 2007).

MI is the ratio between the number of monopolised baits and occurrence, that is the number of visited baits. A bait was considered monopolised by a species when, after 60 min, one of these three conditions occurred:

- the species was the only one at the bait and it occurred with at least 5 individuals (SANTINI *et al.*, 2007);
- there were two species at the bait and one was present with twice as many ants as the other one;
- there were two species at the bait and the non-monopolising one occurred with less than 5 individuals.

DAI is the ratio of discovered baits (i.e. the baits where a species is first to arrive) and occurrence.

The influence of food, time of day and their interactions on species occurrence was tested through a two-way independent ANOVA. Food and time of day were included as fixed factors. To test the existence of competition trade-offs, a Spearman's rank correlation coefficient (rho) was used to test pairwise correlations between the behavioural indices. All analysis were carried out using R version 3.0.2 (CRAN project software).

RESULTS

Baiting permitted to collect 20413 specimens belonging to 3 subfamilies, 14 genera and 23 species (Table 1). In the whole study area, two-way independent ANOVA on occurrence for each species underlined no significant effect for food, time of day nor their interactions (Table 1). Acquiring this result allowed considering the four transects placed in each site as four replicas of the same environment. Therefore, in order to evaluate the existence of a dominance-discovery trade-off in ant communities of Castelporziano, data from all transects of one site were pooled to calculate behavioural indices values for each species. Correlation tests between MI and DAI were not significant considering the whole community (Spearman rank correlation, $r_s = 0.369$, $p = 0.083$). According to high and low vegetation classification, the trade-off was still absent in the first group ($r_s = 0.222$, $p = 0.446$), whereas analyses showed a positive significant correlation for the community of low vegetation sites ($r_s = 0.676$, $p = 0.016$) (Fig. I, 1). In particular, MI and DAI were significantly correlated in WG ($r_s = 1$, $p = 0.040$) (Fig. I, 2) but not in CF ($r_s = 0$, $p = 1$) and in DG ($r_s = -0.389$, $p = 0.611$). Correlation tests for the remaining sites were not significant (PW: $r_s = -0.272$, $p = 0.728$; PIG: $r_s = -0.308$, $p = 0.552$; MOW: $r_s = 0.144$, $p = 0.786$; MS: no monopolising species). Nevertheless, the comparison between behavioural indices of the same species in different sites underlined some interesting differences. *Messor minor* changed indices values from DG to CF (DG: DAI = 0.45, MI = 0.6; CF: DAI = 0, MI = 1) as well as *Tetramorium caespitum* (DG: DAI = 0.5, MI = 0.5; CF: DAI = 1, MI = 1). Moreover, *Plagiolepis pygmaea* showed a reversal in indices values from PW to PIG (PW: DAI = 1, MI = 0; PIG: DAI = 0, MI = 1).

DISCUSSION

According to baiting data, there was no evidence for the existence of a dominance-discovery trade-off in the studied Mediterranean ant communities. On the contrary, a positive correlation between discovery ability (DAI) and monopolisation (MI) was detected in some of the environments chosen for the analyses. Therefore, we agree with PARR and GIBB (2012) suggesting that the dominance-discovery trade-off does not always play a major role in structuring ant assemblages. It is likely that multiple mechanisms contribute to facilitate species coexistence (ANDERSEN, 2008). For example, other trade-offs, such as between thermal tolerance and competitive ability, are likely to play a role in some habitats (CERDÀ *et al.*, 1998, 2013; BESTELMEYER, 2000; WIESCHER *et al.*, 2011). Partitioning the activity space and time is another way that ants use to avoid or reduce competition within their

Table 1 - Species Check list and Two-way independent ANOVA on species occurrence at baits. Food and time of day were included as fixed factors. No significant effect was detected for the factors nor their interactions.

Species	Factor	d.f.	F	P	Species	Factor	d.f.	F	P
<i>Aphaenogaster subterranea</i> (Latreille 1798)	Food	1, 24	2.0833	0.1613	<i>Pheidole pallidula</i> (Nylander 1849)	Food	1, 24	0.1412	0.7104
	Time slot	1, 24	0.0000	1.0000		Time slot	1, 24	0.5647	0.4597
	Interaction	1, 24	0.0000	1.0000		Interaction	1, 24	0.0000	1.0000
<i>Camponotus lateralis</i> (Olivier 1792)	Food	1, 24	1.0000	0.3273	<i>Plagiolepis pygmaea</i> (Latreille 1798)	Food	1, 24	0.3429	0.5637
	Time slot	1, 24	1.0000	0.3273		Time slot	1, 24	0.3429	0.5637
	Interaction	1, 24	1.0000	0.3273		Interaction	1, 24	0.3429	0.5637
<i>Camponotus vagus</i> (Scopoli 1763)	Food	1, 24	0.3333	0.5691	<i>Solenopsis fugax</i> (Latreille 1798)	Food	1, 24	0.6667	0.4222
	Time slot	1, 24	0.3333	0.5691		Time slot	1, 24	0.6667	0.4222
	Interaction	1, 24	0.3333	0.5691		Interaction	1, 24	0.0000	1.0000
<i>Crematogaster scutellaris</i> (Olivier 1792)	Food	1, 24	1.7363	0.2000	<i>Tapinoma subboreale</i> Seifert 2012	Food	1, 24	0.0105	0.9191
	Time slot	1, 24	0.5034	0.4848		Time slot	1, 24	0.8526	0.3650
	Interaction	1, 24	0.8322	0.3707		Interaction	1, 24	0.0947	0.7609
<i>Formica cunicularia</i> Latreille 1798	Food	1, 24	0.2658	0.6109	<i>Tapinoma erraticum</i> (Latreille 1798)	Food	1, 24	1.0000	0.3273
	Time slot	1, 24	0.4394	0.5137		Time slot	1, 24	1.0000	0.3273
	Interaction	1, 24	0.0488	0.8270		Interaction	1, 24	1.0000	0.3273
<i>Lasius lasiooides</i> (Emery 1869)	Food	1, 24	1.0000	0.3273	<i>Temnothorax alienus</i> Schulz, Heinze & Pusch, 2007	Food	1, 24	0.0000	1.0000
	Time slot	1, 24	1.0000	0.3273		Time slot	1, 24	0.0000	1.0000
	Interaction	1, 24	1.0000	0.3273		Interaction	1, 24	0.0000	1.0000
<i>Lasius paralienus</i> Seifert 1992	Food	1, 24	0.1935	0.6639	<i>Temnothorax nylanderi</i> (Foerster, 1850)	Food	1, 24	1.0000	0.3273
	Time slot	1, 24	0.0000	1.0000		Time slot	1, 24	1.0000	0.3273
	Interaction	1, 24	0.7742	0.3876		Interaction	1, 24	1.0000	0.3273
<i>Messor capitatus</i> (Latreille 1798)	Food	1, 24	2.0769	0.1625	<i>Temnothorax parvulus</i> (Schenck, 1852)	Food	1, 24	0.5647	0.4597
	Time slot	1, 24	2.0769	0.1625		Time slot	1, 24	2.2588	0.1459
	Interaction	1, 24	2.0769	0.1625		Interaction	1, 24	0.5647	0.4597
<i>Messor minor</i> (André 1883)	Food	1, 24	0.1200	0.7320	<i>Temnothorax unifasciatus</i> (Latreille, 1798)	Food	1, 24	0.9494	0.3396
	Time slot	1, 24	0.1200	0.7320		Time slot	1, 24	0.9494	0.3396
	Interaction	1, 24	0.0000	1.0000		Interaction	1, 24	0.3418	0.5643
<i>Messor wasmanni</i> Krausse 1910	Food	1, 24	0.2222	0.6416	<i>Tetramorium caespitum</i> (Linnaeus, 1758)	Food	1, 24	0.2182	0.6446
	Time slot	1, 24	0.2222	0.6416		Time slot	1, 24	0.0000	1.0000
	Interaction	1, 24	0.0000	1.0000		Interaction	1, 24	0.8727	0.3595
<i>Monomorium monomorium</i> Bolton 1987	Food	1, 24	0.0681	0.7963	<i>Tetramorium moravicum</i> Novák & Sadil, 1941	Food	1, 24	0.4000	0.5331
	Time slot	1, 24	0.2207	0.6427		Time slot	1, 24	1.6000	0.2180
	Interaction	1, 24	0.0245	0.8769		Interaction	1, 24	0.4000	0.5331
<i>Myrmica scabrinodis</i> (Nylander 1846)	Food	1, 24	0.7826	0.3851					
	Time slot	1, 24	0.0870	0.7706					
	Interaction	1, 24	0.0000	1.0000					




Fig. I - Correlation analyses between the behavioural indices MI and DAI. In pooled low vegetation sites (1), a positive significant correlation was detected. (1): $r_s = 0.676$, $p = 0.016$. (2): $r_s = 0.832$, $p = 0.040$.

community (CROS *et al.*, 1997; ALBRECHT and GOTELLI, 2001; MOONEY and TILLBERG, 2005; SOLIDA *et al.*, 2011, 2014).

Moreover, habitat complexity and vegetative structure are among the main factors affecting community dynamics (CERDÀ *et al.*, 2013). For example, the increased heterogeneity and rugosity of structurally complex habitats may reduce the chance of species interactions, thus facilitating coexistence (PARR and GIBB, 2012). It was likely the case of Wet Grassland (WG), according to the results we found in our previous study with pitfall traps (SPOTTI *et al.*, 2010) and the present one with baits. Maybe the higher level of complexity and diversification of this habitat supported a variegated ant assemblage, as we found with pitfall trapping, even if there were species extremely dominant from a behavioural point of view.

In addition, the present results highlighted a trend of some species behaviour to be affected by the site type. Actually, we recorded a change in monopolisation and discovery ability of the same species in different sites. These differences could not be detected considering only data on numerical dominance, but they add important pieces of information on community dynamics. Further investigations are needed to understand what are the main factors that may influence this aspect, such as vegetative structure or competitive interactions. We conclude that an integrated approach considering numerical and behavioural data is a useful tool to analyse ant community coexistence mechanisms more in details.

ACKNOWLEDGEMENTS

This research has been supported by grants (FIL - 2007) from Italian Ministry of Education, University and Research (MIUR) assigned to Pr. A. Mori and by grants (2007) from Accademia Nazionale delle Scienze detta dei XL assigned to Dr. Cristina Castracani. We wish to thank Dr. Fabrizio Rigato (Museum of Natural History, Milan, Italy) for the identification of specimens. We are grateful to the General Secretariat of Presidency of the Italian Republic, to the Direction of Castelporziano Presidential Estate and to the Observatory of the Multidisciplinary Centre of Mediterranean Costal Ecosystems for the hospitality and the possibility to set up the research.

REFERENCES

- AA.VV. (Eds), 2001 – *Il sistema ambientale della Tenuta Presidenziale di Castelporziano. Ricerche sulla complessità di un ecosistema forestale costiero mediterraneo.* - Accademia Nazionale delle Scienze detta dei XL, Scritti e Documenti, 26, 2 volumi, 866 pp.
- AA.VV. (Eds), 2006 – *Il sistema ambientale della Tenuta Presidenziale di Castelporziano. Ricerche sulla complessità di un ecosistema forestale costiero mediterraneo.* - Accademia Nazionale delle Scienze detta dei XL, Scritti e Documenti, 37, 3 volumi, 1853 pp.
- ADLER R.F., LEBRUN E.G., FEENER JR D.H., 2007 – *Maintaining diversity in an ant community: modeling, extending and testing the dominance-discovery trade-off.* – Am. Nat., 169: 323–333.
- AGOSTI D., MAJER J., ALONSO L., SCHULTZ T., 2000 – *Ants: standard methods for measuring and monitoring biodiversity.* – Smithsonian Institution Press, Washington DC, 280 pp.
- ALBRECHT M., GOTELLI N.J., 2001 – *Spatial and temporal niche partitioning in grassland ants.* – Oecologia, 126: 134–141.
- ANDERSEN A.N., 1992 – *Regulation of “momentary” diversity by dominant species in exceptionally rich ant communities of the Australian seasonal tropics.* – Am. Nat., 140: 401–420.
- ANDERSEN A.N., 2008 – *Not enough niches: non-equilibrium processes promoting species coexistence in diverse ant communities.* – Aust. Ecol. 33: 211–220.
- ARNAN X., GAUCHEREL C., ANDERSEN A.N., 2011 – *Dominance and species co-occurrence in highly diverse ant communities: a test of the interstitial hypothesis and discovery of a three-tiered competition cascade.* – Oecologia, 166: 783–794.
- BESTELMEYER B.T., 2000 – *The trade-off between thermal tolerance and behavioural dominance in a subtropical South American ant community.* – J. Anim. Ecol., 69: 998–1009.
- BOLTON B., 1994 – *Identification guide to the ant genera of the world.* – Harvard University Press, 222 pp.
- CASTRACANI C., GRASSO D.A., FANFANI A., MORI A., 2010 – *The ant fauna of Castelporziano Presidential Reserve (Rome, Italy) as a model for the analysis of ant community structure in relation to environmental variation in Mediterranean ecosystems.* – J. Insect Conserv., 14: 585–594.
- CERDÀ X., ARNAN X., RETANA J., 2013 – *Is competition a significant hallmark of ant (Hymenoptera: Formicidae) ecology?* – Myrmecol. News, 18: 131–147.
- CERDÀ X., RETANA J., CROS S., 1997 – *Thermal disruption of transitive hierarchies in Mediterranean ant communities.* – J. Anim. Ecol., 66: 363–374.
- CERDÀ X., RETANA J., CROS S., 1998 – *Critical thermal limits in Mediterranean ant species: trade-off between mortality risk and foraging performance.* – Funct. Ecol., 12: 45–55.
- CROS S., CERDÀ X., RETANA J., 1997 – *Spatial and temporal variations in the activity patterns of Mediterranean ant communities.* – Ecoscience, 4: 269–278.
- DAVIDSON D.W., 1998 – *Resource discovery versus resource domination in ants: a functional mechanism for breaking the trade-off.* – Ecol. Entomol., 23: 484–490.
- DELLA ROCCA A.B., PIGNATTI S., MUGNOLI S., BIANCO P.M., 2001 – *La carta della vegetazione della Tenuta di Castelporziano. In: Il sistema ambientale della Tenuta Presidenziale di Castelporziano. Ricerche sulla complessità di un ecosistema forestale costiero mediterraneo,* AA.VV. Eds, Accademia Nazionale delle Scienze detta dei XL, Scritti e Documenti, 26, pp. 709–747.
- DELSINNE T., ROISIN Y., LEPONCE M., 2007 – *Spatial and temporal foraging overlaps in a Chacoan ground-foraging ant assemblage.* – J. Arid Environ., 71: 29–44.
- FEENER JR D.H., 2000 – *Is the assembly of ant communities mediated by parasitoids?* – Oikos, 90: 79–88.
- FEENER D.H., ORR M.R., WACKFORD K.M., LONGO J.M., BENSON W.W., GILBERT L.E., 2008 – *Geographic variation in resource dominance-discovery in Brazilian ant communities.* – Ecology, 89: 1824–1836.
- FELLERS J.H., 1987 – *Interference and exploitation in a guild of woodland ants.* – Ecology, 68: 1466–1478.
- GIBB H., PARR C.L., 2010 – *How does habitat complexity affect foraging success? A test using functional measures on three continents.* – Oecologia, 164: 1061–1073.
- HERBERS J.M., 1989 – *Community structure in north temperate ants: temporal and spatial variation.* – Oecologia, 81: 201–211.

- HÖLLODOBLER B., WILSON E.O., 1990 – *The Ants*. – Belknap, Cambridge, Massachusetts, 746 pp.
- HOLWAY D.A., 1999 – *Competitive mechanisms underlying the displacement of native ants by the invasive Argentine ant*. – *Ecology*, 80: 238–251.
- LEBRUN E.G., FEENER JR D.H., 2007 – *When trade-offs interact: balance of terror enforces dominance-discovery trade-off in a local ant assemblage*. – *J. Anim. Ecol.*, 76: 58–64.
- LESSARD J.P., DUNN R.R., SANDERS N.J., 2009 – *Temperature-mediated coexistence in temperate forest ant communities*. – *Insect. Soc.*, 56: 149–156.
- MOONEY K.A., TILLBERG C.V., 2005 – *Temporal and spatial variation to ant omnivory in pine forests*. – *Ecology*, 86: 1225–1235.
- PARR C.L., SINCLAIR B.J., ANDERSEN A.N., GASTON K.J., CHOWN S.L., 2005 – *Constraints and competition in assemblages: a cross-continental and modelling approach for ants*. – *Am. Nat.*, 165: 481–494.
- PARR C.L., GIBB H., 2010 – *Competition and the role of dominant ants*. In: *Ant ecology*, Lach L., Parr C.L., Abbott K.L. Eds., Oxford University Press, New York, pp. 77–96.
- PARR C.L., GIBB H., 2012 – *The discovery–dominance trade-off is the exception, rather than the rule*. – *J. Anim. Ecol.*, 81: 233–241.
- PEARCE-DUVET J.M.C., FEENER JR D.H., 2010 – *Resource discovery in ant communities: do food type and quantity matter?* – *Ecol. Entomol.*, 35: 549–556.
- PERFECTO I., 1994 – *Foraging behavior as determinant of asymmetric competitive interaction between two ant species in a tropical agroecosystem*. – *Oecologia*, 98: 184–192.
- SANDERS N.J., GORDON D.M., 2003 – *Resource-dependent interactions and the organization of desert ant communities*. – *Ecology*, 84: 1024–1031.
- SANTINI G., TUCCI L., OTTONETTI L., FRIZZI F., 2007 – *Competition trade-offs in the organisation of a Mediterranean ant assemblage*. – *Ecol. Entomol.*, 32: 319–326.
- SAVOLAINEN R., VEPSÄLÄINEN K., 1988 – *A competition hierarchy among boreal ants: impact on resource partitioning and community structure*. – *Oikos*, 51: 135–155.
- SOLIDA L., CELANT A., LUISELLI L., GRASSO D.A., MORI A., FANFANI A., 2011 – *Competition for foraging resources and coexistence of two syntopic species of Messor harvester ants in Mediterranean grassland*. – *Ecol. Entomol.*, 36: 409–416.
- SOLIDA L., LUISELLI L., GRASSO D.A., D'EUSTACCHIO D., MORI A., FANFANI A., 2014 – *Spatio-temporal foraging dynamics in two coexisting harvester ants (Hymenoptera: Formicidae)*. – *J. Insect. Behav.*, 27: 429–438.
- SPOTTI F.A., CASTRACANI C., GRASSO D.A., FANFANI A., MORI A., 2010 – *The community structure temporal development of Castelporziano ant fauna*. – *Redia*, XCIII: 89–93.
- STUBLE K.L., RODRIGUEZ-CABAL M.A., MCCORMICK G.L., JURIC I., DUNN R.R., SANDERS N.J., 2013 – *Tradeoffs, competition, and coexistence in eastern deciduous forest ant communities*. – *Oecologia*, 171: 981–992.
- WIESCHER P.T., PEARCE-DUVET J.M.C., FEENER JR D.H., 2011 – *Environmental context alters ecological trade-offs controlling ant coexistence in a spatially heterogeneous region*. – *Ecol. Entomol.*, 36: 549–559.