

FABRIZIO PENNACCHIO (*) - LEONARDO MARIANELLI (*) - FRANCESCO BINAZZI (*)
 VALERIA FRANCARDI (*) - FRANCESCO PAOLI (*) - RAFFAELE GRIFFO (**) - PIO FEDERICO ROVERSI (*)

FIRST INTERCEPTION OF *TRICHOFERUS CAMPESTRIS* (FALDERMANN, 1835) (COLEOPTERA CERAMBYCIDAE CERAMBYCINAE) IN ITALY

(*) Consiglio per la ricerca in agricoltura e l'analisi dell'economia agraria - Research Centre for Agrobiology and Pedology, via Lanchiola 12/a, 50125, Firenze, Italy; E-mail: fabrizio.pennacchio@crea.gov.it

(**) Servizio Fitosanitario Regione Campania. Centro Direzionale - Isola A6 - 80146, Napoli.

Pennacchio F., Marianelli L., Binazzi F., Francardi V., Paoli F., Griffi R., Roversi P.F. – First interception of *Trichoferus campestris* (Faldemann, 1835) (Coleoptera Cerambycidae Cerambycinae) in Italy.

The Velvet Longhorned Beetle *Trichoferus campestris* (Faldemann, 1835) was intercepted for the first time in Italy, in the Naples harbour, during monitoring activities at entry points carried out in the context of the national project ASPROPI. *T. campestris* is widely polyphagous and able to colonize several woody species in both agricultural and forest environments. Moreover timber and solid wood items can be equally attractive to this pest. This additional report, in an area where other alien insect species had been identified in the past, emphasises how the national monitoring network represents a key element of the alert system designed to quickly counter the accidental introduction and spreading of exotic insect pests.

KEY WORDS: Velvet Longhorned Beetle, exotic species, Southern Italy.

INTRODUCTION

Wood is widely used in trades either as a commodity itself or as packaging material for several products. Regrettably, in the absence of phytosanitary measures, it may represent an efficient vector of many xylophagous insects, increasing the risk of pest introductions in new areas (CAVEY, 1998; HAACK, 2001, 2006; HAACK *et al.*, 2010; LIEBOLD *et al.*, 1995 MCCULLOUGH *et al.*, 2006; AUKEEMA *et al.*, 2010; HAACK and RABAGLIA, 2013; PENNACCHIO *et al.*, 2003, 2004, 2013). In recent decades, during monitoring activities, hundreds of xylophagous species, belonging mainly to the families Cerambycidae, Buprestidae, Bostrichidae and Curculionidae (particularly Scolytinae) have been intercepted worldwide in high-risk areas such as ports, airports, customs and nurseries. Since then, a vast number of introduced species have become naturalized showing a strong capacity to spread to new territories, quickly adapting to different ecological conditions. This often led to negative consequences in terms of ecosystem efficiency and host plant health (INGHILESI *et al.*, 2013; PENNACCHIO *et al.*, 2013; HERARD *et al.*, 2009; BULLAS-APPLETON *et al.*, 2014). In this regard, for example, negative outcomes due to the introduction of *Anoplophora chinensis* (Forster, 1771), *A. glabripennis* (Motschulsky, 1853) (Coleoptera Cerambycidae) and *Xylosandrus compactus* (Eichhoff, 1875) (Coleoptera Curculionidae Scolytinae), responsible for severe damages to broad-leaved trees in urban and suburban areas, have been often observed. Moreover *Megaplatypus mutatus* (Chapuis, 1865) and *Aclees* sp. cf. *foveatus* Voss (Coleoptera Curculionidae) have been also reported to be harmful to poplar plantations and fig trees, respectively (TREMBLAY *et al.*, 2000; CIAMPOLINI *et al.*, 2005; MASPERO *et al.*, 2007; ALLEGRO and GRIFFO, 2008; HERARD *et al.*, 2009; PENNACCHIO *et al.*, 2012).

Thanks to the monitoring activities carried out in the project ASPROPI, a specimen of *Trichoferus campestris* (Faldemann, 1835) (Coleoptera Cerambycidae) (Velvet longhorned beetle) was captured by funnel traps located in the port of Naples, Southern Italy: the identification of the specimen was made using the keys of HEGYESSY and KUTASI (2010).

DISTRIBUTION

The genus *Trichoferus* Wollaston, 1854, belongs to the family Cerambycidae and includes 7 species in continental Europe: *T. fasciculatus* (Faldemann, 1837), *T. griseus* (Fabricius, 1792)(Fig. I, 2), *T. holosericeus* (Rossi, 1790) (Fig. I, 3), *T. magnanii* Sama, 1992, *T. pallidus* (Olivier, 1790)(Fig. I, 1), *T. spartii* (G. Müller, 1948) and *T. campestris*. In Italy only *T. fasciculatus*, *T. griseus*, *T. spartii*, *T. pallidus*, *T. arenbergeri* Holzschuh, 1995, endemic of Sardinia and *T. holosericeus* have been recorded (SAMA and RAPUZZI, 2011; RAPUZZI and GREGO, 2013).

T. campestris is native to China, Japan, Korea, Mongolia, India, Kyrgyzstan, Tajikistan, Turkmenistan and Uzbekistan, the easternmost part of Russia, European and south eastern Russia and the southern Urals (KRIVOSHEINA and TOKGAEV, 1985; EPPO, 2009; GREBENNICKOV *et al.*, 2010).

Either stabilized populations or simple interceptions of this species have been recorded in many areas outside its native range such as central and southern Russia, Ukraine, Georgia, Iran, Iraq, Poland, Moldavia, Romania, Moravia (Czech Republic), Slovakia, Hungary, Romania, France, the eastern Mediterranean area, North America, Indian Ocean and Pacific Ocean (DANILEVSKY and MIROSHNIKOV, 1985; MAKHNOVSKII, 1966; DASCĂLU *et al.*, 2013; COQUEMPOT, 2006; SABOL, 2009; SAMA *et al.*, 2005; KRUSZELNICKI, 2010; HEGYESSY and KUTASI, 2010; ZAMOROKA and PANIN, 2011; BULLAS-APPLETON *et al.*, 2014; WATSON *et al.*, 2015; THE




Fig. I – 1. *Trichoferus pallidus* ; 2. *T. griseus* ; 3. *T. holosericeus*; 4. *T. campestris*.

FOOD & ENVIRONMENT RESEARCH AGENCY, 2014). This species has been also intercepted in Israel, Jordan, Lebanon, Syria, northeast Africa (Egypt, Sinai Peninsula), Sweden, Turkey and France (COCQUEMPT, 2006; SABOL, 2009; BOZKURT *et al.*, 2013; DASCĂLU *et al.*, 2013; THE FOOD & ENVIRONMENT RESEARCH AGENCY, 2014).

BIOLOGY

T. campestris (Fig. I, 4) develops in woody plants in both agricultural and forest environments. It is active during the night, flying regularly from the end of June to the beginning of August when females lay eggs into the bark of tree trunks and large-diameter branches. Healthy or stressed plants are usually colonized but any wood material, though very dehydrated, can be equally attractive. In laboratory conditions, *T. campestris* is able to colonize plants and to develop even in small-diameter branches from 2.5 to 3.0 cm. Young larvae initially form galleries between the tree bark and the cambium and then penetrate into the xylem. In fact, bark was often reported to play a key role in *T.*

campestris oviposition and subsequent development of its larvae, particularly in the first phase of their life (IVATA and YAMADA, 1990). On the other hand, older larvae usually feed on the external xylem showing a strong capacity to adapt to their substrate even when it is strongly dehydrated. This species usually overwinters in the larval stage under the bark or in pupal chambers and its galleries can be up to 12 mm broad. Some authors argued that its life cycle is completed in two years (PLAVILSHCHIKOV, 1940; PAVLOVSKII and STACKELBERG, 1955; MAKHNOVSKIY, 1966; KOSTIN, 1973) while others stressed the importance to complete the cycle in a period longer than two years (CHEREPANOV, 1981; DANILEVSKII and MIROSHNIKOV, 1985; LER, 1996).

HOST PLANTS

T. campestris is widely polyphagous and able to colonize several woody species representing essential components of forests, orchards, and street tree populations in Italy and more generally in the EPPO countries. It has been observed to colonize healthy plants belonging to several species included in the genera: *Picea*, *Pinus*, *Betula*, *Broussonetia*, *Gleditsia*, *Malus*, *Morus*, *Salix*, *Sorbus*; when the wood is dehydrated, it is able to exploit an increased number of plant genera and species including: *Abies* spp., *Larix* spp., *Picea* spp., several *Pinus* such as *Pinus densiflora* and also *Chamaecyparis obtusa*, *Betula*, *Fagus crenata*, *Juglans mandshurica*, *Morus bombycina*, *Robinia pseudoacacia*, *Zelkova serrata*, *Vitis vinifera*, *Acer* spp., *Alnus* spp., *Aralia* spp., *Camellia japonica*, *Carpinus*, *Citrus*, *Cornus*, *Diospyros*, *Euonymus*, *Fraxinus*, *Ilex*, *Malus*, *Populus*, *Pyrus*, *Rhus*, *Salix*, *Syzygium*, *Tilia*, *Ulmus*, *Wisteria*, *Quercus*, *Zanthoxylum*, *Ziziphus* (ORLINSKI, 2006; EWBBB, 2014).

PHYTOSANITARY RISK

In 2007, *T. campestris* was included in the EPPO A2 action list and it is currently considered a quarantine insect not only in the USA and Canada but also in the EPPO countries where there is a risk of introduction. In fact, together with its congeneric species reported in the European fauna, it is often responsible for damages to live trees, dehydrated wood, and, in some circumstances, solid wood items. Therefore, since this species has the potential to establish permanently in the vast majority of regions of central and southern Europe, including Italy, it is regarded as a possible threat to biodiversity (MAKHNOVSKII, 1966; KOSTIN, 1973; KRIVOSHEINA and TOKGAEV, 1985).

PHYTOSANITARY MEASURES AIMED AT LIMITING PEST SPREADING

The main vector, responsible for the artificial introduction of *T. campestris* in new geographic areas, is certainly infested wood and in particular packaging material that may host larvae, pupae and adults.

In 2002, specimens of *T. campestris* emerged in a quarantine area of the Marseille harbour (France) from *Salix* sp. timber shipped from China (COCQUEMPT, 2006). In 2015, in Austria, *T. campestris* specimens were intercepted 3 times on packaging material from China marked with the ISPM-31 (EUROPHYT, 2016). Thanks to these last interceptions, made also with the help of dogs, live larvae, pupae and adults of this species could be detected.

Since this pest is able to develop even in highly dehydrated wood, trade of packaging material represents indeed the main spreading route. So far, phytosanitary measures adopted to limit pest introductions included the

early debarking of logs and accurate wood inspections aimed at detecting characteristics diagnostic of xylophagous insects (i.e. frass and gallery systems).

However, we stress that the compliance with the current international legislation concerning phytosanitary measures for packaging material, in agreement with the FAO requirements (ISPM No. 15) (2013), is essential in the implementation of strategies aimed at preventing unintentional pest transfer.

This specimen was collected thanks to a national monitoring network established in the Naples harbour by MIPAAF in cooperation with the regional phytosanitary services, CREA and other scientific institutions. In the same area, an outbreak of *Aromia bungii* (Faldermann, 1835) (Coleoptera Cerambycidae) had been detected in 2012 (EPPO, 2012; EPPO, 2015).

Such results underline that the active monitoring of areas at risk of pest introduction, represents a key element of the alert system designed to quickly counter the accidental introduction and spreading of exotic insects in the Italian territory.

ACKNOWLEDGMENTS

This work was carried out in the context of the National Project “ASPROPRI - Activities for the protection of plants”, funded by MIPAAF (Italian Ministry of Agricultural Food and Forestry Policies, DM n. 28635, 2015/12/28).

We wish to thank the phytosanitary inspector Dr. Elio Minucci (Port of Naples), who has been actively involved in monitoring.

REFERENCES

ALLEGRO G., GRIFFO R., 2008 – *I rischi di diffusione di Megaplatypus mutatus*. - Informatore Agrario, 64 (13): 73-76.

AUKEMA J.E., MCCULLOUGH D.G., VON HOLLE B., LIEBOLD A.M., BRITTON K., FRANKEL, S.J., 2010 – *Historical accumulation of nonindigenous forest pests in the continental US*. - BioScience, 60, 886–897.

BOZKURT V., ÖZDEM A., AYAN E., 2013 – *Coleopteran pests intercepted on imported forest products in Turkey*. Book of proceedings of IV International Symposium “Agrosym 2013”. Jahorina, October 3 - 6, 2013: 646-652.

BULLAS-APPLETON E., KIMOTO T., TURGEON J.J., 2014 – *Discovery of Trichoferus campestris (Coleoptera: Cerambycidae) in Ontario, Canada and first host record in North America*. - Can. Entomol. 146: 111-116.

CAVEY, J.F., 1998 – *Solid wood packing material from China, initial pest risk assessment on certain wood boring beetles known to be associated with cargo shipments: Asian Longhorned Beetle (Anoplophora glabripennis), Ceresium, Monochamus and Hesperophanes*. USDA PPQ, 22pp.

CHEREPANOV A.I., 1981 – *Longhorn Beetles of Northern Asia (Cerambycidae)*. Nauka Press, Novosibirsk, Russia. 216 pp (in Russian). English translation: *Cerambycidae of Northern Asia*. Volume 2, Part 1 Cerambycinae. E.J. Brill, Leiden, The Netherlands. 292 pp.

CIAMPOLINI M., PERRIN H., REGALIN R., 2005 – *Aclees cibratus, nuovo per l’italia nocivo al fico allevato in vivaio*. - Informatore Agrario, 61: 69-71.

COCQUEMPOT, C., 2006 – *Alien longhorned beetles (Coleoptera Cerambycidae): Original interceptions and introductions in Europe, mainly in France, and notes about recently imported species*. - Redia, 89: 35-50.

DANILEVSKY, M.L., MIROSHNIKOV A.I., 1985 – *Longhorn beetles of Caucasus (Coleoptera, Cerambycidae)*. *Taxonomic Keys*. - Nauka Press, Krasnodar, Russia. 419 pp (in Russian).

DASCĂLU, M.M., SERAFIM, R., LINDELOW, A., 2013 – *Range expansion of Trichoferus campestris (Faldermann) (Coleoptera: Cerambycidae) in Europe with the confirmation of its presence in Romania*. - Entomologica Fennica, 24: 142-146.

EPPO, 2009 – *Hesperophanes campestris*. - Bulletin OEPP/EPPO 39: 51-54.

EPPO 2012 – *First report of Aromia bungii in Italy*. - EPPO Reporting Service, 2012-10-01, 2012/204.

EPPO, 2015 - *Aromia bungii*. EPPO Bull, 45: 4-8. doi:10.1111/epb.12173

EUROPHYT, 2016 – data processed - <https://webgate.ec.europa.eu/europhyt>.

EWBBB_2014, Exotic Wood Borer/Bark Beetle - *Trichoferus campestris* (Faldermann), datasheet [https://www.ippc.int/static/media/files/publication/en/2016/01/ISPM_15_2013_En_2015-12-22_PostCPM10_InkAmReformatted.pdf](http://www.google.it/url?sa=t&rct=j&q=&esrc=s&source=web&cd=5&cad=rja&uact=8&ved=0ahUKEwjBp7f9NPLAhXKDywKHRQeDeQQFghCMAQ&url=http%3A%2F%2Fdownload.ceris.purdue.edu%2Ffile%2F2204&u sg=AFQjCNGovtT5xpn_m_oMMH8VAxUboPDzQ&si g2=nxD8AV9E5Pk40JCgUqYu5w&bvm=bv.117218890, d.bGg. (last date access, 22th march 2016)</p>
<p>FAO, 2013 – <i>ISPM 15, Guidelines for regulating wood packaging material in international trade</i>. <a href=).

GREBENNIKOV, V.V., GILL B.D., VIGNEAULT R., 2010 – *Trichoferus campestris (Faldermann) (Coleoptera: Cerambycidae), an Asian wood-boring beetle recorded in North America*. - The Coleopterists Bulletin 64: 13-20.

HAACK R.A., 2001 – *Intercepted Scolytidae (Coleoptera) at U.S. ports of entry: 1985–2000*. - Integrated Pest Management Reviews, 6: 253-282.

HAACK R.A., 2006 – *Exotic bark- and wood-boring Coleoptera in the United States: recent establishments and interceptions*. - Can. J. For. Res., 36: 269-288.

HAACK R.A., HÉRARD F., SUN J., TURGEON J.J., 2010 – *Managing invasive population of Asian Longhorned Beetle and Citrus Longhorned Beetle: a worldwide perspective*. - Ann. Rev. Ent., 55: 521-546.

HAACK R.A., RABAGLIA R.J., 2013 – *Exotic bark and ambrosia beetles (Coleoptera: Curculionidae: Scolytinae) in the United States: potential and current invaders*. In: Peña JE (ed) *Potential invasive pests of agricultural crops*. CAB International, Wallingford, UK, pp. 48-74.

HEGYESSY G., KUTASI C.S., 2010 – *Trichoferus species new to Hungary (Coleoptera: Cerambycidae)*. - Folia Entomologica Hungarica, 71: 35-41.

HERARD F., MASPERO M., RAMUALDE N., JUCKER C., COLOMBO M., CIAMPITI M., CAVAGNA B., 2009 – *Anoplophora glabripennis infestation (Col.: Cerambycidae) in Italy*. - Bulletin OEPP/EPPO Bulletin 39: 146-152.

INGHILESI A.F., MAZZA G., CERVO R., GHERARDI F., SPOSIMO P., TRICARICO E., ZAPPAROLI M., 2013 – *Alien insects in Italy: Comparing patterns from the regional to European level*. - Journal of Insect Science, 13:73, 13 pp., <http://www.insectscience.org/13.73>.

IVATA R., YAMADA F., 1990 – *Notes on the biology of Hesperophanes campestris (Faldermann) (Col.*,

Cerambycidae), a drywood borer in Japan. - Material and Organismen, 25(4): 305-313.

KOSTIN I.A., 1973 - [Dendrophagous Beetles of Kazakhstan (Buprestidae, Cerambycidae, Ipidae)]. Izdatel'stvo Instituta Zoologii Akademii Nauk Kazakhskoi SSR, Alma-Ata, 278 p. (in Russian).

KRIVOSHEINA N.P., TOKGAEV T.V., 1985 - [The formation of trunk insect complexes on irrigated lands in the Kopet-Dag foothills]. Izvestiya Akademii Nauk Turkmeneskoi SSR (5), 34-39, Biologicheskie Nauki, No. 5 (in Russian).

KRUSZELNICKI L., 2010 - Trichoferus campestris (Faldermann, 1835) (Coleoptera: Cerambycidae) in Poland. Acta entomologica silesiana, 18: 33-34 (in Polish).

LER P.A. (Ed.), 1996 - [Taxonomic Keys for Insects of Russian Far East. v. III, part 3, Coleopterous, or Beetles]. Dal'nauka, Vladivostok, 556 p. (in Russian).

LIEBOLD A.M., MACDONALD W.L., BERGDAHL D., MASTRO V.C., 1995 - Invasion by Exotic Forest Pests: A Threat to Forest Ecosystems. - Forest Science Monographs 30, 49 pp.

MAKHNOVSKII I.K., 1966 - [Mulberry longhorn beetle Trichoferus campestris Fald]. In: Vrediteli Gornykh Lesov I Bor'ba S Nimi, pp. 88-89. Lesnaya Promyshlennost', Moscow, (in Russian).

MASPERO, M., CAVALIERI, G., D'ANGELO, G., JUCKER, C., VALENTINI, M., COLOMBO M., HERARD F., LOPEZ J., RAMUALDE N., CIAMPITI M., CAREMI G., CAVAGNA B., 2007 - Anoplophora chinensis - eradication program in Lombardia (Italy). https://www.eppo.int/QUARANTINE/special_topics/anoplophora_chinensis/chinensis_IT_2007.htm

MCCULLOUGH D.G., WORK T.T., CAVEY J.F., LIEBOLD A.M., MARSHALL D., 2006 - Interceptions of nonindigenous plant pests at US ports of entry and border crossings - sings over a 17-year period. - Biol. Invasions, 8: 611-630.

ORLINSKI, A. D., 2006 - Outcomes of the EPPO project on quarantine pests for forestry. - EPPO Bulletin, 36: 497-511.

PAVLOVSKII E.N., SHTAKELBERG A.A. (Ed.), 1955 - [Forest pests. Guide]. Izdatel'stvo Zoologicheskogo Instituta Akademii Nauk SSSR, Moscow - Leningrad, 1, 421 pp. (in Russian).

PENNACCHIO F., ROVERSI P.F., FRANCARDI V., GATTI E., 2003 - Xylosandrus crassiusculus (Motschulsky) a bark beetle new to Europe. - Redia, 86: 77-80.

PENNACCHIO F., FAGGI M., GATTI E., CARONNI F., COLOMBO M., ROVERSI P.F., 2004 - First record of Phloeotribus liminaris (Harris) in Europe (Coleoptera Scolytidae). - Redia, 87: 85-89.

PENNACCHIO F., SANTINI L., FRANCARDI V., 2012 - Bioecological notes on Xylosandrus compactus (Eichhoff) (Coleoptera Curculionidae Scolytinae), a species recently recorded into Italy. - Redia, 95: 67-77.

PENNACCHIO F., DANTI R., BENASSAI D., SQUARCINI M., MARZIALI L., DI LONARDO V., ROVERSI P.F., 2013 - A new additional record of Phloeosinus armatus Reitter from Italy (Coleoptera Curculionidae Scolytinae). - Redia, 96: 45-50.

PLAVILSHCHIKOV N.N., 1940 - [Trichoferus campestris (Fald.)]. In: Fauna SSSR, Coleopterous insects, V. XXII; Longhorn beetles, Part 2, pp. 69-71. Izdatel'stvo Akademii Nauk SSSR, Moskva-Leningrad (RU) (in Russian).

RAPUZZI P. and GREGO B., 2013. First record of Trichoferus pallidus (Olivier, 1790) for Italy (Coleoptera: Cerambycidae). Munis Entomology & Zoology, 8 (2): 712-713.

SABOL, O., 2009 - Trichoferus campestris (Coleoptera: Cerambycidae) - a new species of longhorn beetle for the Czech Republic and Slovakia. - Klapalekiana, 45: 199-201.

SAMA, G., FALLANDHZADEH M., RAPUZZI P., 2005 - Notes on some Cerambycidae (Coleoptera) from Iran with description of two new species (Insecta Coleoptera Cerambycidae). - Quaderno di Studi e Notizie di Storia Naturale della Romagna, 20: 123-132.

SAMA G., RAPUZZI P., 2011 - Una nuova Checklist dei Cerambycidae d'Italia. - Quad. Studi Nat. Romagna, 32: 121-164.

THE FOOD & ENVIRONMENT RESEARCH AGENCY; 2014 - Rapid Pest Risk Analysis for Trichoferus campestris. 7 pp. <https://secure.fera.defra.gov.uk/phiw/riskRegister/plant-health/documents/trichoferusCampestris.pdf>

TREMBLAY, E., ESPINOSA B., MANZINI D., CAPRIO G.. 2000 - Un coleottero proveniente dal Sudamerica minaccia i pioppi. - L'Informatore Agrario 48: 89-90.

WATSON K., PRATT C.A., CAPUTO J., 2015 - Total Records of Velvet Longhorn Beetle Trichoferus campestris Faldermann (Coleoptera, Cerambycidae) from Utah. http://ag.utah.gov/documents/Insect_Velvet_Longhorn_Beetle.pdf.

ZAMOROKA, A. M., PANIN, R. Y., 2011 - Recent records of rare and new for Ukrainian Carpathians species of Longhorn beetles (Insecta: Coleoptera: Cerambycidae) with notes on their distribution. - Munis Entomology & Zoology, 6 (1): 155-165.